

Levi-Civita Connections on the Quantum Groups $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$

István Heckenberger, Konrad Schmüdgen, Universität Leipzig, Fakultät für Mathematik und Informatik und NTZ, Augustusplatz 10, 04109 Leipzig, Germany.

e-mail: istvan@aix550.informatik.uni-leipzig.de, schmuedgen@mathematik.uni-leipzig.d400.de

Abstract

For bicovariant differential calculi on quantum groups various notions on connections and metrics (bicovariant connections, invariant metrics, the compatibility of a connection with a metric, Levi-Civita connections) are introduced and studied. It is proved that for the bicovariant differential calculi on $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$ from the classification in [18] there exist unique Levi-Civita connections.

0 Introduction

The seminal work of S. L. Woronowicz [20] was the starting point to study non-commutative bicovariant differential calculi on quantum groups (Hopf algebras). Woronowicz has developed a general theory of such calculi which in many aspects can be considered as a non-commutative version of the classical Lie group theory. Bicovariant differential calculi on the quantum matrix groups $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$ have been classified (under natural assumptions) in two recent papers [18] and [19]. An outcome of this classification is that except for finitely many values of q there are precisely $2N$ such calculi on $SL_q(N)$ for $N \geq 3$ and two on $O_q(N)$ and $Sp_q(N)$ for $N \geq 4$. See Section 1 for a brief review. It is clear that these calculi are basic tools of non-commutative geometry on the corresponding quantum groups.

The aim of this paper is to define and to study invariant metrics and Levi-Civita connections for the bicovariant differential calculi on $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$. The Killing metric of a compact Lie group and the Levi-Civita connection of a Riemannian manifold are fundamental notions of (commutative) differential geometry, so it seems that extensions of these concepts to the bicovariant differential calculi are necessary steps toward the development of non-commutative differential geometry on quantum groups. It turns out that these generalizations are by no means straightforward and that phenomena occur which are absent in classical differential geometry. We briefly mention some of these new features. Firstly, no ad-invariant metric for these bicovariant differential calculi is symmetric in the usual sense if q is not a root of unity. However, the ad-invariant metrics are symmetric with respect to the corresponding braiding, see Corollary 2.5 below. In our opinion, this fits nicely into the concepts of the braided geometry, see [15] and the references therein. Secondly, ad-invariant metrics for $SL_q(N)$ resp. $O_q(N)$, $Sp_q(N)$ depend on two resp. three complex parameters. Roughly speaking, this stems from the fact that the dimension of these

bicovariant calculi is N^2 rather than the dimension of the corresponding Lie groups as in case of the classical differential calculus. Thirdly, if we generalize the notion of a Levi-Civita connection in obvious straightforward manner, then there exist many Levi-Civita connections for a given metric. For our bicovariant differential calculi they depend on a number of free complex parameters, see e. g. appendix B.

The main purpose of this paper is to propose notions of compatibility of a connection and an ad-invariant metric (see formula (9) and Definition 3.3 below) and of a Levi-Civita connection (see Definition 3.5) which ensure that the important result in classical differential geometry of uniqueness of the Levi-Civita connection remains valid in the present setting. If the braiding map σ is the flip operator of the tensor product as in case of the “ordinary” differential calculus on compact Lie groups, our definitions give just the corresponding classical concepts.

This paper is organized as follows. In Section 1 we briefly recall the bicovariant differential calculi on $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$ studied in [18] and we collect some basic facts needed later. In Section 2 we define metrics and invariant metrics. The invariant metrics for these calculi and their restrictions to the invariant subspaces for the right coaction and the right adjoint action, respectively, are determined. For some of the calculi the quantum Lie algebra contains the corresponding classical Lie algebra as ad-invariant subspace when $q \rightarrow 1$. In these cases the limits of the invariant metrics exist and their restrictions to these subspaces give multiples of the Killing forms. In Appendix A we show how a variant of the Rosso form of the quantized enveloping algebra $\mathcal{U}_q(sl(N))$ gives an ad-invariant metric for $SL_q(N)$. Section 3 is concerned with connections. After reviewing some generally known definitions on connections (see e. g. [4]) we define bicovariant connections, the compatibility of a connection with a metric and Levi-Civita connections and we discuss some of their properties. The main results of this paper (Theorems 4.2 and 5.2) are stated and proved in Sections 4 and 5, respectively. They assert that for each of the bicovariant differential calculi on $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$ described in Section 1 there is precisely one Levi-Civita connection. Moreover, it is shown that these Levi-Civita connections admit limits when $q \rightarrow 1$ in an appropriate way. In appendix B of this paper we show that if we define Levi-Civita connections by taking the “usual” compatibility with a metric, then the set of Levi-Civita connections for $SL_q(N)$ depends on three free parameters.

Let us fix some notation and assumptions which are needed in the sequel. Throughout we use Sweedler’s notation $\Delta(a) = a_{(1)} \otimes a_{(2)}$ and the Einstein convention to sum over repeated indices. The antipode of a Hopf algebra is denoted by κ and the counit by ε . Let $\text{Mor}(v, w)$ the space of intertwiners of corepresentations v and w and let $\text{Mor}(v) := \text{Mor}(v, v)$. We use the definitions of the quantum groups $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$ and their basic properties established in [8]. Let $u = (u_j^i)$ denote the corresponding fundamental representation and let u^c be the contragredient representation of u . In case where $\mathcal{A}$ is $O_q(N)$ or $Sp_q(N)$ let $C = (C_j^i)$ denote the matrix of the metric of $\mathcal{A}$ and B the inverse matrix of C . We abbreviate $Q = q - q^{-1}$.

Also we shall freely use the general theory of bicovariant differential calculi developed in [20]. The abbreviation FODC means a first order differential calculus. As noted above, our main intention is to study the bicovariant differential calculi $\Gamma_{\pm, z}$ (see Section 1) which occurred in the classification of [18, 19]. But the corresponding concepts and general facts apply to an arbitrary bicovariant differential calculus over a Hopf algebra. In order to avoid confusion, let us adopt the following

notations: $\tilde{\Gamma}$ is always a general bicovariant FODC over a Hopf algebra $\mathcal{A}$, $\mathcal{A}^0$ is the Hopf dual of $\mathcal{A}$, $\tilde{\mathcal{X}}$ is the quantum Lie algebra of $\tilde{\Gamma}$, $\{\eta_i | i \in \mathcal{I}\}$ is a finite vector space basis of ${}_{\text{inv}}\tilde{\Gamma}$, $\{\chi_i | i \in \mathcal{I}\}$ is the corresponding dual basis of $\tilde{\mathcal{X}}$ and $\tilde{\sigma}$ is the braiding of $\tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma}$ as defined by Proposition 3.1 in [20].

In this paper we suppose that the deformation parameter q is **not a root of unity** and $q \neq 0$. Then, roughly speaking, the representation theory of $\mathcal{A}$ is similar to the classical case [13, 17]. We shall need this assumption only in order to ensure that the decompositions of certain tensor product representations of u and u^c can be labelled by Young tableaux similar to the classical case. The corresponding results are the Lemmas 2.3, 2.4, 4.1 and 5.1. All other considerations are valid without this assumption.

1 Review of some facts on bicovariant differential calculi on quantum groups $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$

Firstly we repeat the construction of bicovariant differential calculi, see [18] for some missing details of proofs in the following discussion.

Let z be a nonzero complex number. We assume that $z^N = q$ for $\mathcal{A} = SL_q(N)$ and that $z^2 = 1$ for $\mathcal{A} = O_q(N)$ and $\mathcal{A} = Sp_q(N)$. Let $L_z^\pm = (l_{zj}^{\pm i})$ denote the $N \times N$ matrix of linear functionals $l_{zj}^{\pm i}$ on $\mathcal{A}$ defined in [8], Section 2, by taking the matrix $z^{-1}P\hat{R}$ as R . By definition, we then have

$$l_{zj}^{+i}(u_m^n) = z^{-1}\hat{R}_{mj}^{in} \quad \text{and} \quad l_{zj}^{-i}(u_m^n) = z\hat{R}_{mj}^{-1in}. \quad (1)$$

Let $D_i := q^{2i}$ for $\mathcal{A} = SL_q(N)$ and $D_i := (C(C^{-1})^t)_i^i$ for $\mathcal{A} = O_q(N)$, $Sp_q(N)$, where C is the matrix given by the metric of $O_q(N)$ and $Sp_q(N)$, cf. [8], Sect. 1. Then we have $\kappa^2(u_j^i) = D_i u_j^i D_j^{-1}$ (no summation).

There are $2N$ bicovariant FODC $\Gamma_{\pm,z}$, $z^N = q^2$ on $SL_q(N)$, $N \geq 3$, and 2 bicovariant FODC $\Gamma_+ = \Gamma_{+,1}$ and $\Gamma_- = \Gamma_{+,-1}$ on $O_q(N)$, $Sp_q(N)$ and $SL_q(2)$. Except for the quantum group $O_q(3)$, these FODC exhaust the bicovariant first order differential calculi which appeared in the classification of [18, 19]. They are the objects of our study in the paper.

In what follows we assume that Γ always denotes such a FODC $\Gamma_{\pm,z}$. The FODC $\Gamma_{\pm,z}$ on $\mathcal{A}$ is given by

$$da = \sum_{i,j=1}^N (\chi_{ij} * a) \eta_{ij}, \quad a \in \mathcal{A},$$

where

$$\chi_{ij} = \sum_{n=1}^N D_n^{-1} l_{z'i}^{\pm n} \kappa(l_{z''n}^{\mp j}) - D_i^{-1} \delta_{ij} \varepsilon, \quad i, j = 1, \dots, N \quad (2)$$

and $\{\eta_{ij} | i, j = 1, \dots, N\}$ is a basis of the space ${}_{\text{inv}}(\Gamma_{\pm,z})$ of left-invariant elements of $\Gamma_{\pm,z}$. The right and left $\mathcal{A}$ -module operations satisfy the equations $\eta_{ij}a = (f_{mn}^{ij} * a) \eta_{mn}$, $a \in \mathcal{A}$, and the right coaction of $\mathcal{A}$ on this basis is given by $\Delta_R(\eta_{ij}) = \eta_{mn} \otimes v_{ij}^{mn}$, $i, j = 1, \dots, N$, where

$$f_{mn}^{ij} = l_{z'm}^{\pm i} \kappa(l_{z''j}^{\mp n}), \quad (3)$$

$$v_{ij}^{mn} = (u^c u)_{ij}^{mn} = \kappa(u_m^i) u_j^n. \quad (4)$$

The linear span $\mathcal{X}$ of the linear functionals χ_{ij} , $i, j = 1, \dots, N$, equipped with the bracket $[\cdot, \cdot] : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{X}$ defined in [20], Section 5, is called the **quantum Lie algebra** of the bicovariant FODC $\Gamma_{\pm, z}$.

There is a duality $\langle \cdot, \cdot \rangle$ between ${}_{\text{inv}}\tilde{\Gamma}$ and $\tilde{\mathcal{X}}$ given by

$$\left\langle \sum_i a_i \eta_i, \sum_j \chi_j b_j \right\rangle = \sum_k a_k b_k \quad (5)$$

for all $a_k, b_k \in \mathbb{C}$. This definition extends to a map $\langle \cdot, \cdot \rangle : A \otimes {}_{\text{inv}}\tilde{\Gamma} \times \tilde{\mathcal{X}} \otimes B \rightarrow A \otimes_{\mathcal{A}} B$ for linear subspaces A of $\tilde{\Gamma}^\wedge$ resp. $\tilde{\mathcal{X}} \otimes \mathcal{A}$ and linear subspaces B of $\tilde{\Gamma}^\wedge$ by taking $a_k \in A$ and $b_k \in B$ in (5) for $k \in \mathcal{I}$.

Let σ denote the braiding map of $\Gamma_{\pm, z} \otimes_{\mathcal{A}} \Gamma_{\pm, z}$ (see Proposition 3.1 in [20]) and let σ_{ijkl}^{mnrs} be the matrix coefficients of σ with respect to the basis $\{\eta_{ij} \otimes_{\mathcal{A}} \eta_{kl}\}$ of ${}_{\text{inv}}(\Gamma_{\pm, z} \otimes_{\mathcal{A}} \Gamma_{\pm, z})$, i. e. $\sigma(\eta_{ij} \otimes \eta_{kl}) = \sigma_{ijkl}^{mnrs} \eta_{mn} \otimes \eta_{rs}$.

LEMMA 1.1. $\sigma_{ijkl}^{mnrs} = D_k D_x^{-1} \hat{R}_{xj}^{\mp pk} \hat{R}_{yr}^{\pm tn} \hat{R}_{tm}^{\mp xi} \hat{R}_{pl}^{\pm ys}$, where the upper and lower signs refer to $\Gamma_{+, z}$ and $\Gamma_{-, z}$, respectively.

PROOF : We carry out the proof for $\Gamma_{+, z}$. From (3) we see that $f_{rs}^{tp} = l_{z'r}^{+t} \kappa(l_{z''p}^{-s})$ are the linear functionals of [20], Theorem 2.1, for the FODC $\Gamma_{+, z}$. (1) yields $f_{rs}^{tp}(u_l^n) = z^{-1} \hat{R}_{yr}^{tn} \hat{R}_{pl}^{ys}$ and so $\delta_r^i \delta_s^j \delta_m^n = f_{rs}^{ij}(\kappa(\kappa(u_k^n) u_m^k)) = f_{tp}^{ij}(\kappa(u_m^k)) f_{rs}^{tp}(\kappa^2(u_k^n))$. From $\kappa^2(u_k^n) = D_n D_k^{-1} u_k^n$ we easily derive $\delta_r^i \delta_s^j \delta_m^n = f_{tp}^{ij}(\kappa(u_m^k)) z^{-1} D_n D_k^{-1} \hat{R}_{xr}^{tn} \hat{R}_{pk}^{xs}$, so the latter yields $f_{tp}^{ij}(\kappa(u_m^k)) = z D_k D_x^{-1} \hat{R}_{xj}^{-1pk} \hat{R}_{tm}^{-1xi}$. By the general theory (cf. formula (3.15) in [20]) and (4), we have $\sigma_{ijkl}^{mnrs} = f_{rs}^{ij}((u^c)_k^m u_l^n) = f_{tp}^{ij}(\kappa(u_m^k)) f_{rs}^{tp}(u_l^n)$ from which the above formula follows. ■

Recall that the matrix $\hat{R}$ satisfies the Hecke relation $(\hat{R} - qI)(\hat{R} + q^{-1}I) = 0$ for $\mathcal{A} = SL_q(N)$ and the cubic equation $(\hat{R} - qI)(\hat{R} + q^{-1}I)(\hat{R} - \epsilon q^{\epsilon-N}I) = 0$ for $\mathcal{A} = O_q(N)$, $Sp_q(N)$, where $\epsilon = 1$ for $\mathcal{A} = O_q(N)$ and $\epsilon = -1$ for $\mathcal{A} = Sp_q(N)$. From these equations and Lemma 1.1 it follows that

$$(\sigma - I)(\sigma + q^{-2}I)(\sigma + q^2I) = 0$$

for $\mathcal{A} = SL_q(N)$ and

$$(\sigma - I)(\sigma + q^{-2}I)(\sigma + q^2I)(\sigma - \epsilon q^{N-\epsilon+1}I)(\sigma - \epsilon q^{\epsilon-N-1}I)(\sigma + \epsilon q^{N-\epsilon-1}I)(\sigma + \epsilon q^{\epsilon-N+1}I) = 0$$

for $\mathcal{A} = O_q(N)$, $Sp_q(N)$ (cf. [3]).

The above formulas show that the set of for $q > 0$ positive eigenvalues of σ is $\{1\}$ for $SL_q(N)$ and $\{1, q^N, q^{-N}\}$ for $O_q(N)$ and $Sp_q(N)$.

In the following we use the abbreviations $p := \epsilon q^{N-\epsilon}$, $\mathfrak{s} := 1 + Q^{-1}(p - p^{-1})$ for $O_q(N)$ and $Sp_q(N)$ and $\mathfrak{s} := \sum_{i=1}^N q^{-2i}$ for $SL_q(N)$.

Let $\omega_{ij} := \kappa(u_n^i) du_j^n$, $i, j = 1, \dots, N$. In [18] and [19] it is proved that if q is not a root of unity and apart from finitely many other values of q the set $\{\omega_{ij} \mid i, j = 1, \dots, N\}$ is a basis of the vector space ${}_{\text{inv}}\Gamma$. Let $\{X_{ij} \mid i, j = 1, \dots, N\}$ be the corresponding dual basis of $\mathcal{X}$ with respect to the duality (5). We call the sets $\{\omega_{ij}\}$ and $\{X_{ij}\}$ **standard bases** of ${}_{\text{inv}}\Gamma$ and $\mathcal{X}$, respectively. For this basis of ${}_{\text{inv}}\Gamma$ the right coaction also fulfills $\Delta_R(\omega_{ij}) = \omega_{kl} \otimes v_{ij}^{kl}$ (cf. (4)).

Let us briefly return to a general bicovariant FODC $\tilde{\Gamma}$ over a Hopf algebra $\mathcal{A}$ such that $\dim_{\text{inv}} \tilde{\Gamma} < \infty$. Then the quantum Lie algebra $\tilde{\mathcal{X}}$ of $\tilde{\Gamma}$ is contained in the Hopf dual $\mathcal{A}^0$ and the bracket $[\cdot, \cdot]$ of $\tilde{\mathcal{X}}$ can be written as $[x, y] = \text{ad}_R y(x)$. Here ad_R denotes the right adjoint action of $\mathcal{A}^0$ given by $\text{ad}_R f(g) = \kappa(f_{(1)})g f_{(2)}$ for $f, g \in \mathcal{A}^0$. Moreover, we have $[x, f] := \text{ad}_R f(x) \in \tilde{\mathcal{X}}$ for all $f \in \mathcal{A}^0$ and $x \in \tilde{\mathcal{X}}$. A linear subspace $\mathcal{Y}$ of $\tilde{\mathcal{X}}$ is called ad-invariant if $[y, f] \in \mathcal{Y}$ for all $y \in \mathcal{Y}$ and $f \in \mathcal{A}^0$.

Next we are looking for ad-invariant subspaces of $\tilde{\mathcal{X}}$. For this we need the following simple

LEMMA 1.2. *Suppose that $A = (A_j^i) \in \text{Mor}(\tilde{v})$, i. e. $A_j^i \tilde{v}_k^j = \tilde{v}_l^i A_k^l$ for all $i, k \in \mathcal{I}$. Then $\text{im } A = \text{lin}\{A_j^i \eta_i \mid j \in \mathcal{I}\}$ is a Δ_R -invariant subspace of $\text{inv} \tilde{\Gamma}$ and $\text{im } A^t = \text{lin}\{A_j^i \chi_j \mid i \in \mathcal{I}\}$ is an ad-invariant subspace of $\tilde{\mathcal{X}}$.*

PROOF : Since $A \in \text{Mor}(\tilde{v})$, we have $\Delta_R(A_j^i \eta_i) = A_j^i \eta_k \otimes \tilde{v}_i^k = A_l^k \eta_k \otimes \tilde{v}_j^l$, so that $\Delta_R(\text{im } A) \subset \text{im } A \otimes \mathcal{A}$. From the general theory [20] we easily derive that $[\chi_i, f] = f(\tilde{v}_k^i) \chi_k$ for all $f \in \mathcal{A}^0$. Therefore, we compute $\text{ad}_R f(A_k^i \chi_k) = A_k^i f(\tilde{v}_l^k) \chi_l = f(\tilde{v}_j^i) A_l^j \chi_l$. That is, $\text{ad}_R f(\text{im } A^t) \subset \text{im } A^t$ for all $f \in \mathcal{A}^0$. ■

For $SL_q(N)$ the projections $P_0, P_1 : \text{inv} \Gamma \rightarrow \text{inv} \Gamma$ defined by $P_0^{ij} = \frac{1}{s} q^{-2i} \delta^{ij} \delta_{kl}$, $P_1^{ij} = \delta_k^i \delta_l^j - P_0^{ij}$ belong to $\text{Mor}(u^c \otimes u)$. Hence the subspaces $\Upsilon^0 = \text{im } P_0 = \text{lin}\{\omega^0 := P_0(\omega_{11}) = P_0^{kl} \omega_{kl}\}$ and $\Upsilon^1 = \text{im } P_1 = \text{lin}\{\omega_{ij}^1 := P_1(\omega_{ij}) = P_1^{kl} \omega_{kl}\}$ of $\text{inv} \Gamma$ are Δ_R -invariant. The corresponding ad-invariant subspaces of $\mathcal{X}$ are $\mathcal{Y}^0 = \text{im } P_0^t = \text{lin}\{Y^0 := \frac{1}{s} \sum_k X_{kk}\}$ and $\mathcal{Y}^1 = \text{im } P_1^t = \text{lin}\{Y_{ij}^1 := P_1^{ij} X_{kl} = X_{ij} - q^{-2i} \delta^{ij} Y^0\}$ respectively. Since $\{\omega_{ij}\}$ and $\{X_{ij}\}$ are dual bases, we also have $da = \sum_{ij} (X_{ij} * a) \omega_{ij}$ for all $a \in \mathcal{A}$. Because of $P_0 + P_1 = \text{id}$ and $P_0 P_1 = P_1 P_0 = 0$, the latter leads to the formula

$$da = \sum_{i,j=1}^N (Y_{ij}^1 * a) \omega_{ij}^1 + s(Y^0 * a) \omega^0, \quad a \in \mathcal{A}.$$

Now we turn to the quantum groups $O_q(N)$ and $Sp_q(N)$. Then the projections $P_0, P_+, P_- : \text{inv} \Gamma \rightarrow \text{inv} \Gamma$ given by

$$P_0^{ij} = \frac{1}{s} B_m^{ti} C^{mj} \delta_{kl}, \quad P_+^{ij} = \frac{1}{q + q^{-1}} (q^{-1} \delta_k^i \delta_l^j + B_m^{ti} \hat{R}_{nl}^{mj} C_k^{tn} - (q^{-1} + p^{-1}) P_0^{ij}),$$

$$P_-^{ij} = \frac{1}{q + q^{-1}} (q \delta_k^i \delta_l^j - B_m^{ti} \hat{R}_{nl}^{mj} C_k^{tn} + (p^{-1} - q) P_0^{ij})$$

belong to $\text{Mor}(u^c \otimes u)$. Let denote $P_1 := P_-, P_2 := P_+$ for $O_q(N)$ and $P_1 := P_+, P_2 := P_-$ for $Sp_q(N)$. Then, by Lemma 1.2, the subspaces Υ^0, Υ^1 and Υ^2 of $\text{inv} \Gamma$ spanned by the sets $\{\omega^0 := \frac{1}{s} B_k^{tm} C^{kn} \omega_{mn}\}$, $\{\omega_{ij}^1 := P_1(\omega_{ij}) \mid i, j = 1, \dots, N\}$ and $\{\omega_{ij}^2 := P_2(\omega_{ij}) \mid i, j = 1, \dots, N\}$ respectively, are Δ_R -invariant. Moreover, the subspaces $\mathcal{Y}^0, \mathcal{Y}^1$ and $\mathcal{Y}^2$ of the quantum Lie algebra $\mathcal{X}$ generated by the sets $\{Y^0 := \frac{1}{s} \sum_m X_{mm}\}$, $\{Y_{ij}^1 := P_1^{ij} X_{kl} \mid i, j = 1, \dots, N\}$ and $\{Y_{ij}^2 := P_2^{ij} X_{kl} \mid i, j = 1, \dots, N\}$, respectively, are ad-invariant. Similarly as in case of $SL_q(N)$, we obtain the following formula for the differentiation

$$da = \sum_{i,j=1}^N (Y_{ij}^1 * a) \omega_{ij}^1 + \sum_{i,j=1}^N (Y_{ij}^2 * a) \omega_{ij}^2 + s(Y^0 * a) \omega^0, \quad a \in \mathcal{A}.$$

To investigate the classical limit of the structures appearing in this article we keep the basis $\{\omega_{ij} \mid i, j = 1, \dots, N\}$ fixed.

Firstly let $\mathcal{A} = SL_q(N)$. We always consider the classical limit in the sense that $z \rightarrow 1$ and $q \rightarrow 1$, where z is the N -th root of q^2 . (This is not the only possibility, see [10].) For simplicity we shall write $\lim_{q \rightarrow 1}$ for this classical limit. Then, as shown in [10], all functionals X_{ij} , Y^0 and Y_{ij}^1 have limits when $q \rightarrow 1$. It is easily seen that the 1-forms ω^0 and ω_{ij}^1 and the projections P_0 and P_1 have limits as well for $q \rightarrow 1$. We denote these limits by $\vec{Y}^0$, $\vec{Y}_{ij}^1$, $\vec{\omega}^0$, $\vec{\omega}_{ij}^1$, $\vec{P}_0$ and $\vec{P}_1$, respectively. It was proved in [10] that the functionals $\vec{Y}_{ij}^1$ equipped with the limit of the bracket $[\cdot, \cdot]$ are generators of the Lie algebra $sl(N)$.

Now let $\mathcal{A} = O_q(N)$ or $\mathcal{A} = Sp_q(N)$. As proved in [10], for both calculi Γ_+ and Γ_- all functionals X_{ij} , Y^0 , Y_{ij}^1 and Y_{ij}^2 , $i, j = 1, \dots, N$ admit limits when $q \rightarrow 1$. Also, the 1-forms ω^0 , ω_{ij}^1 , ω_{ij}^2 and the projections P_k , $k = 0, 1, 2$ have limits as $q \rightarrow 1$. We shall use the notations $\vec{Y}^0 := \lim_{q \rightarrow 1} Y^0$, $\vec{Y}_{ij}^1 := \lim_{q \rightarrow 1} Y_{ij}^1$, $\vec{Y}_{ij}^2 := \lim_{q \rightarrow 1} Y_{ij}^2$, $\vec{P}_k := \lim_{q \rightarrow 1} P_k$, $k = 0, 1, 2$, $\vec{\omega}^0 := \lim_{q \rightarrow 1} \omega^0$, $\vec{\omega}_{ij}^1 := \lim_{q \rightarrow 1} \omega_{ij}^1$ and $\vec{\omega}_{ij}^2 := \lim_{q \rightarrow 1} \omega_{ij}^2$ for $i, j = 1, \dots, N$. For the FODC $\Gamma = \Gamma_+$ the functionals $\vec{Y}_{ij}^1$, $i, j = 1, \dots, N$ equipped with the limit of the bracket $[\cdot, \cdot]$ span the Lie algebras $o(N)$ and $sp(N)$, respectively.

2 Metrics

We begin with some definitions for a general bicovariant FODC $\tilde{\Gamma}$ over a Hopf algebra $\mathcal{A}$.

DEFINITION 2.1. A bilinear map $g : \tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma} \rightarrow \mathcal{A}$ is called a **metric** on $\tilde{\Gamma}$ if g is nondegenerate (i.e. $g(\xi \otimes \zeta) = 0$ for all $\zeta \in \tilde{\Gamma}$ implies $\xi = 0$, $g(\xi \otimes \zeta) = 0$ for all $\xi \in \tilde{\Gamma}$ implies $\zeta = 0$) and if

$$g(a\xi \otimes \zeta) = ag(\xi \otimes \zeta) \text{ for any } a \in \mathcal{A}, \quad \xi \text{ and } \zeta \in \tilde{\Gamma}. \quad (6)$$

We call a metric g **symmetric** if $g\tilde{\sigma} = g$.

For the ‘‘ordinary’’ differential calculus on Lie groups the braiding map $\tilde{\sigma}$ is just the flip operator, so we obtain the usual notion of a symmetric metric in this case. From condition (6) in the preceding definition it follows easily that a metric g on $\tilde{\Gamma}$ is already completely determined by the elements $g(\eta_i \otimes \eta_j)$, $i, j \in \mathcal{I}$, of $\mathcal{A}$.

For any metric g on $\tilde{\Gamma}$ we have for all $\xi, \zeta \in \tilde{\Gamma}$ and $a \in \mathcal{A}$,

$$\begin{aligned} (\text{id} \otimes \varepsilon g)(\Delta_L(a\xi \otimes \zeta)) &= a(\text{id} \otimes \varepsilon g)(\Delta_L(\xi \otimes \zeta)), \\ (\varepsilon g \otimes \text{id})(\Delta_R(a\xi \otimes \zeta)) &= a(\varepsilon g \otimes \text{id})(\Delta_R(\xi \otimes \zeta)). \end{aligned} \quad (7)$$

DEFINITION 2.2. A metric g on $\tilde{\Gamma}$ is called **invariant** if for all $\xi, \zeta \in \tilde{\Gamma}$,

$$(\text{id} \otimes \varepsilon g)(\Delta_L(\xi \otimes \zeta)) = g(\xi \otimes \zeta) \quad \text{and} \quad (\varepsilon g \otimes \text{id})(\Delta_R(\xi \otimes \zeta)) = g(\xi \otimes \zeta). \quad (8)$$

By (7), the above definition is compatible with the left $\mathcal{A}$ -module structure of $\tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma}$.

Using the representation theory of quantum groups we now show that the invariant metrics on Γ form a 2-parameter family for $SL_q(N)$ and a 3-parameter family for $O_q(N)$ and $Sp_q(N)$.

LEMMA 2.3. Let $\mathcal{A} = SL_q(N)$ and let Γ be as in Section 1. A metric g on Γ is invariant if and only if with complex parameters α and β such that $\alpha \neq 0$, $\alpha + \mathfrak{s}\beta \neq 0$

$$g(\eta_{ij} \otimes \eta_{kl}) = q^{2j}\alpha\delta_{il}\delta_{jk} + \beta\delta_{ij}\delta_{kl}.$$

PROOF : Suppose g is an invariant metric on Γ . From the first equation in (8) it follows that $g(\xi \otimes \zeta) = \varepsilon g(\xi \otimes \zeta)$ for $\xi, \zeta \in \text{inv}\Gamma$. From (4) and the second equation in (8) we conclude that $g \in \text{Mor}(u^c \otimes u \otimes u^c \otimes u, 1)$. Since q is not a root of unity by assumption, the multiplicities of irreducible components in the decomposition of the tensor product representation $u^c \otimes u \otimes u^c \otimes u$ are the same as in the classical case. Therefore, $\dim \text{Mor}(u^c \otimes u \otimes u^c \otimes u, 1) = 2$. Since $q^{2j} u_n^j \kappa(u_j^r) = q^{2n} \delta_{nr}$ by [8], the transformations $T = (T_{ijkl})$ and $S = (S_{ijkl})$ with $T_{ijkl} := q^{2j} \delta_{il} \delta_{jk}$ and $S_{ijkl} := \delta_{ij} \delta_{kl}$ belong to $\text{Mor}(u^c \otimes u \otimes u^c \otimes u, 1)$. Thus we get $g(\eta_{ij} \otimes \eta_{kl}) = q^{2j} \alpha \delta_{il} \delta_{jk} + \beta \delta_{ij} \delta_{kl}$ for some complex numbers α and β . This map is nondegenerate if and only if $\alpha \neq 0$ and $\alpha + \mathfrak{s}\beta \neq 0$.

Conversely, it is easily seen that the above formula defines an invariant metric g on Γ . $\blacksquare$

LEMMA 2.4. *Let $\mathcal{A} = O_q(N)$ or $\mathcal{A} = Sp_q(N)$ and Γ denote one of the FODC from Section 1. A metric g on Γ is invariant if and only if with complex parameters α, β and γ such that $\alpha + p\beta + \mathfrak{s}\gamma \neq 0, \alpha - q\beta \neq 0, \alpha + q^{-1}\beta \neq 0$*

$$g(\eta_{ij} \otimes \eta_{kl}) = ((\alpha B_{14} B_{23} + \beta B_{12} B_{34} \hat{R}_{23} + \gamma B_{12} B_{34}) C_1^t C_3^t)_{ijkl}.$$

In Lemma 2.4 we used the following notation. Let $C = (C_j^i)$ be the matrix of the metric which occurs in the definition of $O_q(N)$ resp. $Sp_q(N)$ (see [8]) and let $B = (B_j^i)$ its inverse matrix. Set $C^t := (C^{ti}_j) = (C^{ji}) = (C^j_i)$ and $B^t := (B^{ti}_j) = (B_{ji}) = (B^j_i)$. Then the notation in Lemma 2.4 is the usual leg numbering notation, i.e. the equation therein reads as

$$g(\eta_{ij} \otimes \eta_{kl}) = (\alpha B_{ml} B_{jn} + \beta B_{mr} B_{sl} \hat{R}_{jn}^{rs} + \gamma B_{mj} B_{nl}) C_i^{tm} C_k^{tn}.$$

PROOF of Lemma 2.4: The proof is similar to the proof of Lemma 2.3. The decomposition of the tensor product gives now $\dim \text{Mor}(u^c \otimes u \otimes u^c \otimes u, 1) = 3$. The conditions on the coefficients ensure the nondegeneracy of the metric g . $\blacksquare$

Some straightforward computations based on Lemma 1.1 and the particular form of the invariant metrics in Lemma 2.3 and 2.4 prove the following

COROLLARY 2.5. *Any invariant metric on Γ is symmetric.*

Let us consider again a general bicovariant FODC $\tilde{\Gamma}$ over an arbitrary Hopf algebra $\mathcal{A}$.

Let g be a metric on $\tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma}$ and let g_{ij} be the matrix coefficients of g , i.e. $g(\eta_i \otimes \eta_j) = g_{ij}$ with respect to a fixed basis $\{\eta_i\}$ of $\text{inv}\tilde{\Gamma}$. Recall that $\{\chi_i\}$ is the dual basis of $\{\eta_i\}$. We suppose that $\{a_j \mid j \in \mathcal{I}\}$ is a finite subset of $\mathcal{A}$ such that $\chi_i(a_j) = \delta_{ij}$.

DEFINITION 2.6. A map $g^* : \tilde{\mathcal{X}} \otimes \tilde{\mathcal{X}} \rightarrow \mathcal{A}$ such that $g^*(\chi_i \otimes \chi_j) = g^{*ij}$ and $\sum_k g_{ik} g^{*kj} = \delta_i^j$ for all $i, j \in \mathcal{I}$ is called the **dual metric of g** . The metric g^* is called **ad-invariant** if

$$g^*([\chi', f_{(1)}] \otimes [\chi'', f_{(2)}]) = f(1) g^*(\chi' \otimes \chi'') \quad \text{for all } \chi', \chi'' \in \tilde{\mathcal{X}}, f \in \mathcal{A}^0. \quad (9)$$

Note that the nondegeneracy of g corresponds to the nondegeneracy of g^* .

There is an interesting link between invariant and ad-invariant metrics given by

PROPOSITION 2.7. *If g is an invariant metric on $\tilde{\Gamma}$ then the dual metric g^* of g is ad-invariant. Conversely, let $g^* : \tilde{\mathcal{X}} \otimes \tilde{\mathcal{X}} \rightarrow \mathbb{C}$ be an ad-invariant metric and suppose that $\mathcal{A}^0$ separates the points of $\mathcal{A}$. Then g is an invariant metric.*

PROOF : In the proof we take the bases of ${}_{\text{inv}}\tilde{\Gamma}$ and $\tilde{\mathcal{X}}$ described above.

Let g be invariant. From the first equation in (8) it follows that $g(\eta_i \otimes \eta_j) = g_{ij} \in \mathbb{C}$ for all $i, j \in \mathcal{I}$. Since g is nondegenerate, the dual metric g^* is defined and is a map to $\mathbb{C} \cdot 1$. The second equation in (8) means that $g_{ij} \tilde{v}_m^i \tilde{v}_n^j = g_{mn}$ which yields $g^{*ij} \tilde{v}_i^m \tilde{v}_j^n = g^{*mn}$ for the dual metric g^* . Applying functionals $f \in \mathcal{A}^0$ to the last equation we get $g^*(f_{(1)}(\tilde{v}_i^m) \chi_i \otimes f_{(2)}(\tilde{v}_j^n) \chi_j) = f(1)g^*(\chi_m \otimes \chi_n)$. Using $[\chi_i, f] = f(\tilde{v}_k^i) \chi_k$ for all $f \in \mathcal{A}^0$ the latter is equivalent to $g^*([\chi_m, f_{(1)}] \otimes [\chi_n, f_{(2)}]) = f(1)g^*(\chi_m \otimes \chi_n)$ for all $m, n \in \mathcal{I}$ and all $f \in \mathcal{A}^0$, i. e. the dual metric is ad-invariant.

Suppose that $g^* : \tilde{\mathcal{X}} \otimes \tilde{\mathcal{X}} \rightarrow \mathbb{C}$ is ad-invariant. Then the matrix elements of g are also complex numbers. This implies the first equation in (8). Reversing the reasoning from the preceding paragraph, it follows from the ad-invariance of g^* that $f(g^{*ij} \tilde{v}_i^m \tilde{v}_j^n - g^{*mn}) = 0$ for all $m, n \in \mathcal{I}$ and $f \in \mathcal{A}^0$. Since $\mathcal{A}^0$ separates the points of $\mathcal{A}$, we obtain $g^{*ij} \tilde{v}_i^m \tilde{v}_j^n = g^{*mn}$ for all $m, n \in \mathcal{I}$ from which the second equation in (8) follows. This proves the invariance of g . $\blacksquare$

Next we specialize again to the bicovariant FODC $\Gamma = \Gamma_{\pm, z}$ over $\mathcal{A} = SL_q(N)$, $O_q(N)$, $Sp_q(N)$. We compute the dual metrics g^* of the invariant metrics g from Lemmas 2.3 and 2.4 and their restrictions to the ad-invariant subspaces of the quantum Lie algebra $\mathcal{X}$. Let us say that two subspaces Υ and Υ' of ${}_{\text{inv}}\Gamma$ are **mutually orthogonal with respect to a metric g** on ${}_{\text{inv}}\Gamma$ if $g(x \otimes x') = g(x' \otimes x) = 0$ for all $x \in \Upsilon$, $x' \in \Upsilon'$. A similar notion is used for the dual metric g^* on $\mathcal{X}$.

For $SL_q(N)$ we have (see Lemma 2.3) $g(\eta_{ij} \otimes \eta_{kl}) = q^{2j} \alpha \delta_{il} \delta_{jk} + \beta \delta_{ij} \delta_{kl}$ and we get

$$g^*(\chi_{ij} \otimes \chi_{kl}) = \left(q^{-2i} \alpha^{-1} \delta_{il} \delta_{jk} - \frac{\beta}{\alpha(\alpha + \mathfrak{s}\beta)} q^{-2i-2k} \delta_{ij} \delta_{kl} \right) \quad (10)$$

for all $i, j, k, l = 1, \dots, N$. For $O_q(N)$ and $Sp_q(N)$ the dual metric of the metric in Lemma 2.4 is

$$g^*(\chi_{ij} \otimes \chi_{kl}) = \frac{B_m^{ti} B_n^{tk}}{(\alpha - q\beta)(\alpha + q^{-1}\beta)} \left(\alpha C_{ml} C_{jn} - \beta \hat{R}^{-1}{}_{rs}^{jn} C_{mr} C_{sl} + \frac{-Q\alpha\beta - \alpha\gamma + p^{-1}\beta\gamma}{\alpha + p\beta + \mathfrak{s}\gamma} C_{mj} C_{nl} \right)$$

for $i, j, k, l = 1, \dots, N$.

From equations (2) and (1) we get the transformation formula between the bases $\{X_{ij}\}$ and $\{\chi_{ij}\}$. Then we can express the generators of the ad-invariant subspaces in terms of the basis $\{\chi_{ij}\}$. For the quantum group $SL_q(N)$ and the FODC $\Gamma = \Gamma_{\pm, z}$ we obtain the formulas $Y^0 = \mathfrak{s}^{-1} \mu_{\pm, z}^{-1} \sum_k \chi_{kk}$ and $Y_{ij}^1 = \nu_{\pm, z}^{-1} P_{1kl}^{ij} \chi_{kl}$. For the quantum groups $O_q(N)$ and $Sp_q(N)$ and the calculus $\Gamma = \Gamma_{\pm 1}$ we get $Y^0 = \mathfrak{s}^{-1} \mu_{0\pm}^{-1} \sum_m \chi_{mm}$, $Y_{ij}^1 = \mu_{1\pm}^{-1} P_{1kl}^{ij} \chi_{kl}$ and $Y_{ij}^2 = \mu_{2\pm}^{-1} P_{2kl}^{ij} \chi_{kl}$. Here the constants are defined by $\mu_{+, z} = \mathfrak{s}(z^{-1} - 1) + z^{-1} q^{-1} Q$, $\mu_{-, z} = \mathfrak{s}(z - 1) - z q^{-2N-1} Q$, $\nu_{+, z} = z^{-1} q^{-1} Q$, $\nu_{-, z} = -z q^{-2N-1} Q$, $\mu_{0+} = (p - p^{-1})Q$, $\mu_{0-} = (p - p^{-1})Q + 2\mathfrak{s}$, $\mu_{1\pm} = \pm p(1 + q^{-N})Q$ and $\mu_{2\pm} = \pm(p - p^{-1} q^N)Q$.

One easily verifies that $\mathcal{Y}^0$ and $\mathcal{Y}^1$ are orthogonal subspaces of $\mathcal{X}$ with respect to the dual metric g^* . The restrictions of g^* to $\mathcal{Y}^0 \otimes \mathcal{Y}^0$ and $\mathcal{Y}^1 \otimes \mathcal{Y}^1$ are given by

$$g^*(Y^0 \otimes Y^0) = \frac{1}{\mathfrak{s}(\alpha + \mathfrak{s}\beta) \mu_{\pm, z}^2}, \quad g^*(Y_{ij}^1 \otimes Y_{kl}^1) = \frac{1}{\alpha \nu_{\pm, z}^2} \left(q^{-2i} \delta_{il} \delta_{jk} - \frac{q^{-2i-2k}}{\mathfrak{s}} \delta_{ij} \delta_{kl} \right)$$

for all $i, j, k, l = 1, \dots, N$.

Then Υ^0 and Υ^1 are orthogonal subspaces of ${}_{\text{inv}}\Gamma$ with respect to all invariant metrics g . If g has the form as in Lemma 2.3, then we have

$$g(\omega^0 \otimes \omega^0) = \frac{(\alpha + \mathfrak{s}\beta)\mu_{\pm,z}^2}{\mathfrak{s}}, \quad g(\omega_{ij}^1 \otimes \omega_{kl}^1) = \alpha\nu_{\pm,z}^2 \left(q^{2j}\delta_{il}\delta_{jk} - \frac{1}{\mathfrak{s}}\delta_{ij}\delta_{kl} \right)$$

for all $i, j, k, l = 1, \dots, N$.

The ad-invariant subspaces $\mathcal{Y}^0$, $\mathcal{Y}^1$ and $\mathcal{Y}^2$ of the quantum Lie algebras $\mathcal{X}$ of $O_q(N)$ and $Sp_q(N)$ are mutually orthogonal with respect to g^* . The restrictions of g^* to $\mathcal{Y}^i \otimes \mathcal{Y}^i$, $i = 0, 1, 2$ are described by the formulas

$$g^*(Y^0 \otimes Y^0) = \frac{1}{\mu_{0\pm}^2 \mathfrak{s} \alpha_0},$$

$$g^*(Y_{ij}^1 \otimes Y_{kl}^1) = \frac{B_m^{ti} B_n^{tk}}{\mu_{1\pm}^2 (p^{-1}q^N + pq^{-N}) \alpha_1} \left(p^{-1}q^N C^{ml} C^{jn} - \hat{R}_{rs}^{-1jn} C^{mr} C^{sl} + \frac{p^{-1}(1-q^N)}{\mathfrak{s}} C^{mj} C^{nl} \right),$$

$$g^*(Y_{ij}^2 \otimes Y_{kl}^2) = \frac{B_m^{ti} B_n^{tk}}{\mu_{2\pm}^2 (p^{-1}q^N + pq^{-N}) \alpha_2} \left(pq^{-N} C^{ml} C^{jn} - \hat{R}_{rs}^{-1jn} C^{mr} C^{sl} - \frac{p^{-1} + pq^{-N}}{\mathfrak{s}} C^{mj} C^{nl} \right),$$

where we use the abbreviations $\alpha_0 := \alpha + p\beta + \mathfrak{s}\gamma$, $\alpha_1 := \alpha - p^{-1}q^N\beta$, $\alpha_2 := \alpha + pq^{-N}\beta$. Note that all denominators in the above formulas are non-zero, since the metric is nondegenerate and Γ is a FODC as in [18].

The corresponding subspaces Υ^0 , Υ^1 and Υ^2 of ${}_{\text{inv}}\Gamma$ are mutually orthogonal with respect to all invariant metrics g . If g is as in Lemma 2.4, then we have

$$g(\omega^0 \otimes \omega^0) = \frac{\mu_{0\pm}^2 \alpha_0}{\mathfrak{s}},$$

$$g(\omega_{ij}^1 \otimes \omega_{kl}^1) = \frac{\mu_{1\pm}^2 \alpha_1}{p^{-1}q^N + pq^{-N}} \left(pq^{-N} B_{ml} B_{jn} - B_{mr} B_{sl} \hat{R}_{jn}^{rs} + \frac{p(1-q^{-N})}{\mathfrak{s}} B_{mj} B_{nl} \right) C_i^{tm} C_k^{tn},$$

$$g(\omega_{ij}^2 \otimes \omega_{kl}^2) = \frac{\mu_{2\pm}^2 \alpha_2}{p^{-1}q^N + pq^{-N}} \left(p^{-1}q^N B_{ml} B_{jn} + B_{mr} B_{sl} \hat{R}_{jn}^{rs} - \frac{p + p^{-1}q^N}{\mathfrak{s}} B_{mj} B_{nl} \right) C_i^{tm} C_k^{tn}$$

for all $i, j, k, l = 1, \dots, N$.

Now we want to examine the classical limits.

Let $\mathcal{A} = SL_q(N)$ and let g be an invariant metric as described in Lemma 2.3. The complex numbers α and β may of course depend on the parameters q and z . Let us assume that the functions $\alpha = \alpha(q, z)$ and $\beta = \beta(q, z)$ are choosen such that the limits $c_1 := \lim_{q \rightarrow 1} Q^2 \alpha(q, z)$ and $c_0 := \lim_{q \rightarrow 1} Q^4 (\alpha(q, z) + \mathfrak{s}\beta(q, z))$ exist and are non-zero. Then it follows immediately from the existence of the classical limits as discussed at the end of Section 1 that the invariant metric g and its dual metric admit limits when $q \rightarrow 1$ and $z \rightarrow 1$. The restriction of the limit of g^* to the linear functionals $\vec{Y}_{ij}^1$, $i, j = 1, \dots, N$, is just a complex multiple of the Killing form for $sl(N)$. Similar results are valid for the quantum groups $O_q(N)$ and $Sp_q(N)$ and for the FODC Γ_+ if we suppose that for the functionals $\alpha_k = \alpha_k(q)$, $k = 0, 1, 2$ the limits $c_0 := \lim_{q \rightarrow 1} \mu_{0+}^2 \alpha_0(q)$, $c_1 := \lim_{q \rightarrow 1} \mu_{1+}^2 \alpha_1(q)$ and $c_2 := \lim_{q \rightarrow 1} \mu_{2+}^2 \alpha_2(q)$ exist and are non-vanishing.

3 Connections

We begin with some general definitions (cf. [4]) which, of course, apply to any differential calculus $\tilde{\Gamma}^\wedge = \bigoplus_{n=0}^\infty \tilde{\Gamma}^n$ over an arbitrary algebra $\mathcal{A}$.

Let $\mathcal{E}$ be a left $\mathcal{A}$ -module. A **left connection** on $\mathcal{E}$ is a linear map $\nabla : \mathcal{E} \rightarrow \tilde{\Gamma} \otimes_{\mathcal{A}} \mathcal{E}$ such that

$$\nabla(a\zeta) = da \otimes \zeta + a \nabla(\zeta) \quad \text{for all } a \in \mathcal{A} \text{ and } \zeta \in \mathcal{E}. \quad (11)$$

Let $\tilde{\Gamma}\mathcal{E} := \tilde{\Gamma}^\wedge \otimes_{\mathcal{A}} \mathcal{E}$ be the “ $\mathcal{E}$ -valued differential forms”. A connection ∇ on $\mathcal{E}$ admits a unique extension to a linear map $\nabla : \tilde{\Gamma}\mathcal{E} \rightarrow \tilde{\Gamma}\mathcal{E}$ of degree one such that

$$\nabla(\alpha\zeta) = (d\alpha)\zeta + (-1)^n \alpha \nabla(\zeta) \quad \text{for } \alpha \in \tilde{\Gamma}^n \text{ and } \zeta \in \tilde{\Gamma}\mathcal{E}.$$

The mapping $R(\nabla) = \nabla^2 : \mathcal{E} \rightarrow \tilde{\Gamma}^2 \otimes_{\mathcal{A}} \mathcal{E}$ is called the **curvature** of the connection ∇ . Clearly, $R(\nabla)$ is $\mathcal{A}$ -linear, i.e. $R(\nabla)(a\zeta) = aR(\nabla)(\zeta)$ for $a \in \mathcal{A}$ and $\zeta \in \mathcal{E}$.

Similar concepts can be defined for a right $\mathcal{A}$ -module $\mathcal{E}$. A right connection on $\mathcal{E}$ is then a linear map $\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \tilde{\Gamma}$ satisfying $\nabla(\zeta a) = \zeta \otimes da + \nabla(\zeta)a$ for $a \in \mathcal{A}$ and $\zeta \in \mathcal{E}$. It extends uniquely to a linear map $\nabla : \mathcal{E} \otimes_{\mathcal{A}} \tilde{\Gamma}^\wedge \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \tilde{\Gamma}^\wedge$ such that $\nabla(\zeta\alpha) = (-1)^n \zeta d\alpha + \nabla(\zeta)\alpha$, $\alpha \in \tilde{\Gamma}^n$ and $\zeta \in \mathcal{E} \otimes_{\mathcal{A}} \tilde{\Gamma}^n$. The curvature of ∇ is $R(\nabla) := \nabla^2 : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{A}} \tilde{\Gamma}^2$.

If ∇ is a connection on a left $\mathcal{A}$ -module $\mathcal{E}$, then there is a connection ∇^* on the right $\mathcal{A}$ -module $\mathcal{E}^*$, called the **dual connection** of ∇ , defined by

$$\langle \xi, \nabla^*(\zeta) \rangle = d \langle \xi, \zeta \rangle - \langle \nabla(\xi), \zeta \rangle \quad \text{for } \xi \in \mathcal{E}, \quad \zeta \in \mathcal{E}^*.$$

By a **right** (resp. **left**) **connection** on an $\mathcal{A}$ -bimodule $\mathcal{E}$ we mean a connection on $\mathcal{E}$ when $\mathcal{E}$ is considered as a right (resp. left) $\mathcal{A}$ -module.

We now specialize to the case of our main interest where $\mathcal{E} = \tilde{\Gamma}$ is considered as a left $\mathcal{A}$ -module. Suppose that $\nabla : \tilde{\Gamma} \rightarrow \tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma}$ is a (left) connection on $\tilde{\Gamma}$. Then the **torsion** $T(\nabla)$ is defined by $T(\nabla) = d - m\nabla$, where $m : \tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma} \rightarrow \tilde{\Gamma}^2$ denotes the multiplication map, i.e. $m(\xi \otimes \zeta) = \xi \wedge \zeta$ for $\xi, \zeta \in \tilde{\Gamma}$. The torsion $T(\nabla)$ is $\mathcal{A}$ -linear, since $T(\nabla)(a\xi) = d(a\xi) - m\nabla(a\xi) = da \wedge \xi + a d\xi - da \wedge \xi - am\nabla(\xi) = aT(\nabla)(\xi)$ for $a \in \mathcal{A}$, $\xi \in \tilde{\Gamma}$ by the Leibniz rule and (11).

DEFINITION 3.1. The connection ∇ is called **bicovariant** if

$$\Delta_L \nabla = (\text{id} \otimes \nabla) \Delta_L \quad \text{and} \quad \Delta_R \nabla = (\nabla \otimes \text{id}) \Delta_R. \quad (12)$$

A simple computation shows that the set of all bicovariant connections on $\tilde{\Gamma}$ forms a complex affine space $\text{BC}(\tilde{\Gamma})$.

Let $\tilde{\Gamma}^\wedge = \bigoplus_{n=0}^\infty \tilde{\Gamma}^n$ be a bicovariant differential calculus over a Hopf algebra $\mathcal{A}$. Let $\{\eta_i \mid i \in \mathcal{I}\}$ be a basis in ${}_{\text{inv}}\tilde{\Gamma}$, $\Delta_R(\eta_i) = \eta_j \otimes \tilde{v}_i^j$, and let $D(\nabla)_i^{jk}$ denote arbitrary elements of $\mathcal{A}$. Then the map $\nabla : {}_{\text{inv}}\tilde{\Gamma} \rightarrow \tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma}$, defined by $\nabla(\eta_i) = D(\nabla)_i^{jk} \eta_j \otimes \eta_k$ extends uniquely to a connection on $\tilde{\Gamma}$ and any connection on $\tilde{\Gamma}$ is of this form.

LEMMA 3.2. Let $\tilde{v}$ denote the corepresentation of $\mathcal{A}$ defined by Δ_R on ${}_{\text{inv}}\tilde{\Gamma}$. The connection ∇ on $\tilde{\Gamma}$ is bicovariant if and only if $D(\nabla)_i^{jk} \in \mathbb{C}$ for all i, j, k and $D(\nabla) = (D(\nabla)_i^{jk}) \in \text{Mor}(\tilde{v}, \tilde{v} \otimes \tilde{v})$.

PROOF : Let ∇ be a bicovariant connection on $\tilde{\Gamma}$ and $\nabla(\eta_i) = D(\nabla)_i^{jk} \eta_j \otimes \eta_k$, $D(\nabla)_i^{jk} \in \mathcal{A}$. Because of the first formula of (12) we have $\Delta(D(\nabla)_i^{jk}) = 1 \otimes D(\nabla)_i^{jk}$ and so $D(\nabla)_i^{jk} \in \mathbb{C} \cdot 1$. The second equation of (12) tells us that $(\nabla \otimes \text{id})\Delta_R(\eta_i) = \eta_j \otimes \eta_k \otimes D(\nabla)_x^{jk} \tilde{v}_i^x = \Delta_R \nabla(\eta_i) = \eta_j \otimes \eta_k \otimes (\tilde{v} \otimes \tilde{v})_{xy}^{jk} D(\nabla)_i^{xy}$, i. e. $D(\nabla)\tilde{v} = (\tilde{v} \otimes \tilde{v})D(\nabla)$.

Let now $D(\nabla)_i^{jk} \in \mathbb{C}$ and $D(\nabla) \in \text{Mor}(\tilde{v}, \tilde{v} \otimes \tilde{v})$. Then the above equations written in reversed order show that (12) is true for the connection ∇ defined by $\nabla(\eta_i) = D(\nabla)_i^{jk} \eta_j \otimes \eta_k$. ■

Now let g be an invariant metric on $\tilde{\Gamma}$, cf. Definition 2.2. Let us introduce some new concepts.

DEFINITION 3.3. We say that the connection ∇ on $\tilde{\Gamma}$ is **compatible with the metric** g if

$$(\text{id} \otimes g)(\nabla(\xi) \otimes \zeta) + (g \otimes \text{id})(\text{id} \otimes \tilde{\sigma})(\xi \otimes \nabla(\zeta)) - \text{dg}(\xi \otimes \zeta) = 0 \quad \text{for any } \xi \in \tilde{\Gamma}, \zeta \in {}_{\text{inv}}\tilde{\Gamma}.$$

LEMMA 3.4. Let g denote an invariant metric. Then a connection ∇ on $\tilde{\Gamma}$ is compatible with g if and only if

$$(\text{id} \otimes g)(\nabla(\eta_i) \otimes \eta_j) + (g \otimes \text{id})(\eta_i \otimes \tilde{\sigma} \nabla(\eta_j)) = 0 \quad \text{for all } i, j \in \mathcal{I}. \quad (13)$$

PROOF : In both directions we use only that the metric g is left-invariant, i. e. the first equation in (8) is fulfilled. This is equivalent to $g_{ij} = g(\eta_i \otimes \eta_j) \in \mathbb{C}$.

Suppose ∇ is compatible with g . Because of the first formula in (8) and Definition 3.3 equation (13) is valid. Let us now assume equation (13) is fulfilled and introduce arbitrary elements $\xi \otimes \zeta = a_{ij} \eta_i \otimes \eta_j \in \tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma}$, $a_{ij} \in \mathcal{A}$. Then the assertion follows from the computation

$$\begin{aligned} & (\text{id} \otimes g)(\nabla(a_{ij} \eta_i) \otimes \eta_j) + (g \otimes \text{id})(a_{ij} \eta_i \otimes \tilde{\sigma} \nabla(\eta_j)) - \text{dg}(a_{ij} \eta_i \otimes \eta_j) = \\ & = (\text{id} \otimes g)((\text{d}a_{ij} \otimes \eta_i + a_{ij} \nabla \eta_i) \otimes \eta_j) + a_{ij} (g \otimes \text{id})(\eta_i \otimes \tilde{\sigma} \nabla \eta_j) - \\ & - (\text{d}(a_{ij})g(\eta_i \otimes \eta_j) + a_{ij} \text{d}g(\eta_i \otimes \eta_j)) = \\ & = a_{ij} ((\text{id} \otimes g)(\nabla \eta_i \otimes \eta_j) + (g \otimes \text{id})(\eta_i \otimes \tilde{\sigma} \nabla \eta_j)) - a_{ij} \text{d}g(\eta_i \otimes \eta_j) \end{aligned}$$

and the assumption $g_{ij} \in \mathbb{C}$. ■

DEFINITION 3.5. Let g be an invariant metric. A bicovariant connection ∇ on $\tilde{\Gamma}$ is called a **Levi-Civita connection** (with respect to the metric g) if ∇ is compatible with g and has vanishing torsion $T(\nabla)$.

Recall that any basis of the vector space ${}_{\text{inv}}\tilde{\Gamma}$ is a free left $\mathcal{A}$ -module basis of $\tilde{\Gamma}$. Therefore, any connection on $\tilde{\Gamma}$, its curvature and its torsion are uniquely determined by its values on such a basis.

Now we consider the pairing between $\tilde{\Gamma}$ and $\tilde{\mathcal{X}} \otimes \mathcal{A}$ as a right $\mathcal{A}$ -module (see formula (5)). Since for a connection ∇ on $\tilde{\Gamma}$ there is a dual connection ∇^* on $\tilde{\mathcal{X}} \otimes \mathcal{A}$, it is natural to ask what the conditions in Definition 3.5 mean in terms of ∇^* . For this we assume that $\tilde{\Gamma}^2 = (\tilde{\Gamma} \otimes_{\mathcal{A}} \tilde{\Gamma}) / \ker(\tilde{\sigma} - \text{id})$.

Suppose that ∇ is a connection on $\tilde{\Gamma}$ and let ∇^* be its dual connection. For arbitrary $\xi, \zeta \in \tilde{\mathcal{X}}$ we define $\nabla_{\xi}^*(\zeta) = \langle \nabla^*(\zeta), \xi \rangle$ and mean the pairing between $\tilde{\mathcal{X}} \otimes \tilde{\Gamma}$ and $\tilde{\mathcal{X}}$ (cf. (5)). For the basis elements η_i of ${}_{\text{inv}}\tilde{\Gamma}$ we write $\nabla(\eta_i) = \Gamma_i^{jk} \eta_j \otimes \eta_k$ with $\Gamma_i^{jk} \in \mathcal{A}$. Recall that by Lemma 3.2, we have $\Gamma_i^{jk} \in \mathbb{C}$ for all $i, j, k \in \mathcal{I}$. Then the pairing gives $\nabla_{\chi_i}^*(\chi_j) = -\Gamma_k^{ij} \chi_k$ for the elements of

the dual basis. Let η denote the left- and right-invariant element of $\tilde{\Gamma}$ used in the definition of the differentiation, i. e. $da = \eta a - a\eta$ for $a \in \mathcal{A}$. Recall that the torsion is $\mathcal{A}$ -linear. Therefore, because of $d\eta_i = \eta \wedge \eta_i + \eta_i \wedge \eta$, the torsion of ∇ is vanishing if and only if $d\eta_i = m\nabla(\eta_i)$ or equivalently

$$(\text{id} - \tilde{\sigma})(\eta \otimes \eta_i + \eta_i \otimes \eta - \nabla(\eta_i)) = 0 \quad \text{for all } i \in \mathcal{I}.$$

Using the notation above the latter is equivalent to the equation

$$\nabla_{\chi_i}^*(\chi_j) - \tilde{\sigma}_{kl}^{ij} \nabla_{\chi_k}^*(\chi_l) = [\chi_i, \chi_j] \text{ for all } i, j \in \mathcal{I}.$$

Suppose that g is an invariant metric. By dualizing the condition in Lemma 3.4, it follows that ∇ is compatible with the metric g if and only if

$$g^*(\chi_i \otimes \nabla_{\chi_j}^*(\chi_k)) + g^*(\tilde{\sigma}_{mn}^{ij} \nabla_{\chi_m}^*(\chi_n) \otimes \chi_k) = 0 \text{ for all } i, j, k \in \mathcal{I}.$$

The above equations show that our concepts are analogous to the corresponding notions in classical differential geometry.

4 Levi-Civita connections on $SL_q(N)$

In this section we examine the differential calculi $\Gamma_{\pm, z}$ for $\mathcal{A} = SL_q(N)$. After a short lemma we will prove our main results.

LEMMA 4.1. *We have $\dim \text{BC}(\Gamma) = 5$ for $N = 2$ and $\dim \text{BC}(\Gamma) = 6$ for $N \geq 3$.*

PROOF : Since q is not a root of unity, decompositions of tensor product representations of $\mathcal{A} = SL_q(N)$ can be labelled by Young tableaux as in the classical case. Therefore, we obtain $u^c \otimes u = [0] \oplus [2, 1^{N-2}]$ and $u^c \otimes u \otimes u^c \otimes u = 2[0] \oplus k[2, 1^{N-2}] \oplus$ other terms with $k = 3$ for $N = 2$ and $k = 4$ for $N \geq 3$. Then by the general representation theory we conclude that $\dim \text{Mor}(v, v \otimes v) = 5$ for $N = 2$ and $\dim \text{Mor}(v, v \otimes v) = 6$ for $N \geq 3$. By Lemma 3.2, a connection ∇ on $\Gamma_{\pm, z}$ is bicovariant if and only if $D(\nabla) \in \text{Mor}(v, v \otimes v)$, where $v = u^c \otimes u$. Thus the assertion of the lemma follows. ■

Let $\Gamma^\wedge = \bigoplus_{n=0}^\infty \Gamma^n$ be a differential calculus over $\mathcal{A}$ which contains $\Gamma_{\pm, z}$ as its first order differential calculus Γ . As in [20] we suppose that $\Gamma^2 = (\Gamma \otimes_{\mathcal{A}} \Gamma) / \ker(\sigma - \text{id})$. Then we have the following

THEOREM 4.2. *For any invariant metric g on Γ there exists precisely one Levi-Civita connection ∇ . If $g(\eta_{ij} \otimes \eta_{kl}) = q^{2j} \alpha \delta_{il} \delta_{jk} + \beta \delta_{ij} \delta_{kl}$ with $\alpha, \beta \in \mathbb{C}, \alpha \neq 0, \alpha + \mathfrak{s}\beta \neq 0$ then for $\Gamma_{+, z}$ this Levi-Civita connection is given by*

$$\begin{aligned} \nabla(\eta_{ij}) = & \frac{Q}{2} (q^{2p-2a-2c} \hat{R}_{am}^{bn} \hat{R}_{cn}^{dp} \hat{R}_{jp}^{im} \eta_{ab} \otimes \eta_{cd} - q^{-1} \eta_{ia} \otimes \eta_{aj} - \\ & - Q \eta_{ij} \otimes \eta - Q \eta \otimes \eta_{ij} + \delta_{ij} Q (1 + \mathfrak{s} \alpha^{-1} \beta) q^{-2a} \eta_{ab} \otimes \eta_{ba} - \\ & - \delta_{ij} (q^{2N+1} Q^2 + Q \alpha^{-1} \beta) \eta \otimes \eta) \end{aligned}$$

and for $\Gamma_{-, z}$ by

$$\begin{aligned} \nabla(\eta_{ij}) = & \frac{Q}{2} (-q^{2p-2a-2c} \hat{R}_{am}^{bn} \hat{R}_{cn}^{dp} \hat{R}_{jp}^{im} \eta_{ab} \otimes \eta_{cd} + q^{-2N-1} \eta_{ia} \otimes \eta_{aj} + \\ & + \delta_{ij} (2Q + Q \mathfrak{s} \alpha^{-1} \beta) q^{-2a} \eta_{ab} \otimes \eta_{ba} + \delta_{ij} (q^{2N+1} Q^2 - Q \alpha^{-1} \beta) \eta \otimes \eta). \end{aligned}$$

PROOF : By Lemma 2.3, any invariant metric g is of the form $g(\eta_{ij} \otimes \eta_{kl}) = q^{2j}\alpha\delta_{il}\delta_{jk} + \beta\delta_{ij}\delta_{kl}$ with $\alpha, \beta \in \mathbb{C}, \alpha \neq 0, \alpha + \mathfrak{s}\beta \neq 0$. From the proof of Lemma 4.1 we know that a connection ∇ on Γ is bicovariant if and only if there are complex numbers $\lambda_1, \dots, \lambda_6$ such that

$$\nabla(\eta_{ij}) = \sum_{n=1}^6 \lambda_n (A_n)_{ij}^{abcd} \eta_{ab} \otimes \eta_{cd}, \quad (14)$$

where $\{A_1, \dots, A_6\}$ generates the vector space $\text{Mor}(v, v \otimes v)$. For our calculi we have $v = u^c \otimes u$. By explicit decompositions of the tensor product representations $u^c \otimes u$ and $u^c \otimes u \otimes u^c \otimes u$ it can be shown that the following 6 morphisms $A_1, \dots, A_6$ ($A_k = (A_{kij}^{abcd})$) span the vector space $\text{Mor}(u^c \otimes u, u^c \otimes u \otimes u^c \otimes u)$:

$$\begin{aligned} A_1 &= q^{-2a-2c}\delta_{ij}\delta^{ab}\delta^{cd} & A_2 &= q^{-2a}\delta_{ij}\delta^{ad}\delta^{bc} & A_3 &= \delta_i^a\delta_j^b q^{-2c}\delta^{cd} \\ A_4 &= q^{-2a}\delta_{ab}\delta_i^c\delta_j^d & A_5 &= \delta_i^a\delta^{bc}\delta_j^d & A_6 &= q^{2n-2a-2d}\hat{R}_{am}^{bn}\hat{R}_{cn}^{dp}\hat{R}_{jp}^{im}. \end{aligned}$$

For our differential calculus we have $d\eta_{ij} = \eta \wedge \eta_{ij} + \eta_{ij} \wedge \eta$. Since $\Gamma^2 = (\Gamma \otimes_{\mathcal{A}} \Gamma) / \ker(\sigma - \text{id})$, the torsion of ∇ vanishes if and only if we have

$$(\sigma - \text{id})(\eta \otimes \eta_{ij} + \eta_{ij} \otimes \eta - \sum_{k=1}^6 \lambda_k A_{kij}^{mnrs} \eta_{mn} \otimes \eta_{rs}) = 0$$

in the tensor product $\Gamma \otimes_{\mathcal{A}} \Gamma$. Comparing the coefficients of basis elements the latter is equivalent to the equations

$$\lambda_3 = \lambda_4 = 1 + \frac{\lambda_5}{q^{-1}Q} - \frac{(Q^2 + 1)\lambda_6}{Q} \quad \text{for } \Gamma_{+,z}, \quad (15)$$

$$\lambda_3 = \lambda_4 = 1 - \frac{\lambda_5}{q^{-2N-1}Q} + \frac{\lambda_6}{Q} \quad \text{for } \Gamma_{-,z}. \quad (16)$$

Using Lemma 3.4 it follows that ∇ satisfies the compatibility condition with the metric g if and only if the following equations are fulfilled:

$$\begin{aligned} \lambda_4 + Q\lambda_6 &= 0, & \lambda_5 + q^{-1}\lambda_6 &= 0, & \alpha\lambda_2 + (\alpha + \mathfrak{s}\beta)\lambda_3 &= 0, \\ (\alpha + \mathfrak{s}\beta)\lambda_1 + \beta\lambda_2 + \beta\lambda_4 + q^{2N}Q(\beta + qQ\alpha)\lambda_6 &= 0 \end{aligned} \quad (17)$$

for $\Gamma_{+,z}$ and

$$\begin{aligned} \lambda_4 &= 0, & \lambda_5 + q^{-2N-1}\lambda_6 &= 0, \\ (\alpha + \mathfrak{s}\beta)\lambda_1 + \beta\lambda_2 + (q^{2N+1}Q^2\alpha + q^{2N}Q\beta)\lambda_6 &= 0, \\ \alpha\lambda_2 + (\alpha + \mathfrak{s}\beta)\lambda_3 + (-q^{2N+1}Q\alpha + \beta)\lambda_5 + (Q\alpha + q^{-1}\beta)\lambda_6 &= 0 \end{aligned} \quad (18)$$

for $\Gamma_{-,z}$. Some straightforward computations show that the equations (15) and (17) resp. (16) and (18) have unique solutions λ_i for $\Gamma_{+,z}$ resp. $\Gamma_{-,z}$. Inserting these solutions into (14) we obtain the formulas given in the theorem.

It is not difficult to check that the connection defined by the above formulas fulfills all conditions for a Levi-Civita connection. $\blacksquare$

Let us look closer at the Levi-Civita connection on $\Gamma = \Gamma_{+,z}$. Using Theorem 4.2 for the left and right invariant element $\eta \in {}_{\text{inv}}\Gamma$ we compute $\nabla(\eta) = \frac{Q^2}{2\alpha}(\alpha + \mathfrak{s}\beta)(\mathfrak{s}q^{-2a}\eta_{ab} \otimes \eta_{ba} - \eta \otimes \eta)$. Transforming the basis we obtain

$$\nabla(\omega^0) = \frac{\mu_{+,z}}{\mathfrak{s}} \nabla(\eta) = \frac{z^2 q^2 \mu_{+,z}(\alpha + \mathfrak{s}\beta)}{2\alpha} q^{-2a} \omega_{ab}^1 \otimes \omega_{ba}^1. \quad (19)$$

Moreover, $\nabla(\eta_{ij} - \delta_{ij}\mathfrak{s}^{-1}\eta) = \frac{Q}{2}(q^{2n-2a-2d}\hat{R}_{am}^{bn}\hat{R}_{cn}^{-1dp}\hat{R}_{jp}^{im} - q^{-1}\delta_i^a\delta^{bc}\delta_j^d - Q\delta^{ab}\delta_i^c\delta_j^d + Q\mathfrak{s}^{-1}q^{-2a-2c}\delta_{ij}\delta^{ab}\delta^{cd})\eta_{ab} \otimes \eta_{cd}$. Transforming the basis once again we get

$$\begin{aligned} \nabla(\omega_{ij}^1) &= \nu_{+,z} \nabla(\eta_{ij} - \delta_{ij}\mathfrak{s}^{-1}\eta) = \\ &= \frac{1}{2}(zqq^{2n-2a-2d}\hat{R}_{am}^{bn}\hat{R}_{cn}^{-1dp}\hat{R}_{jp}^{im}\omega_{ab}^1 \otimes \omega_{cd}^1 - \\ &\quad - z\omega_{ia}^1 \otimes \omega_{aj}^1 - Q^2\mu_{+,z}^{-1}(\omega_{ij}^1 \otimes \omega^0 + \omega^0 \otimes \omega_{ij}^1)). \end{aligned} \quad (20)$$

What happens with the Levi-Civita connection in the classical limit? As explained in Section 1, we consider the classical limit in the sense that $z \rightarrow 1$ and $q \rightarrow 1$. Retaining the notation introduced in Sections 1 and 2, formulas (19) and (20) show that the limit of the Levi-Civita connection exists and takes the form

$$\nabla^{\text{cl}}(\vec{\omega}^0) = \lim_{q \rightarrow 1} \nabla(\omega^0) = \frac{(N^2 - 1)c_0}{4Nc_1} \vec{\omega}_{ab}^1 \otimes \vec{\omega}_{ba}^1, \quad (21)$$

$$\nabla^{\text{cl}}(\vec{\omega}_{ij}^1) = \frac{1}{2} \left(\vec{\omega}_{aj}^1 \otimes \vec{\omega}_{ia}^1 - \vec{\omega}_{ia}^1 \otimes \vec{\omega}_{aj}^1 - \frac{2N}{N^2 - 1} (\vec{\omega}_{ij}^1 \otimes \vec{\omega}^0 + \vec{\omega}^0 \otimes \vec{\omega}_{ij}^1) \right). \quad (22)$$

5 Levi-Civita connections on $O_q(N)$ and $Sp_q(N)$

Now we turn to the differential calculi Γ_+ and Γ_- on the quantum groups $O_q(N)$ and $Sp_q(N)$. We begin with

LEMMA 5.1. *We have $\dim \text{BC}(\Gamma_{\pm}) = 14$ for $Sp_q(4)$ and $\dim \text{BC}(\Gamma_{\pm}) = 15$ for $O_q(N)$, $N \geq 3$ and $Sp_q(N)$, $N > 4$.*

PROOF : The proof is similar to the proof of Lemma 4.1. In the present cases $\mathcal{A} = O_q(N)$ and $\mathcal{A} = Sp_q(N)$ we obtain the decompositions of the tensor product representations $u^c \otimes u = [0] \oplus [2] \oplus [1, 1]$, $u^c \otimes u \otimes u^c \otimes u = 3[0] \oplus 6[2] \oplus k[1, 1] \oplus \text{other terms}$, where $k = 5$ for $Sp_q(4)$ and $k = 6$ for $O_q(N)$, $N \geq 3$ and $Sp_q(N)$, $N > 4$. $\blacksquare$

Let $\Gamma^\wedge = \bigoplus_{n=0}^\infty \Gamma^n$ be a differential calculus over $\mathcal{A}$ such that $\Gamma = \Gamma^1 = \Gamma_{\pm}$. In contrast to the $SL_q(N)$ case we assume that $\Gamma^2 = (\Gamma \otimes_{\mathcal{A}} \Gamma) / \mathcal{K}$, where $\mathcal{K} = \ker(\sigma - \text{id}) \oplus \ker(\sigma - q^N \text{id}) \oplus \ker(\sigma - q^{-N} \text{id})$ (see the last remarks in Section 1). Such an assumption for the higher order calculus has already been used in [3].

In case of the ‘‘ordinary’’ classical differential calculus the 2-forms are the quotient of the tensor product of 1-forms by the eigenspace of the flip operator with eigenvalue 1. Since the eigenvalues

q^N and q^{-N} of the braiding map σ tend to 1 when $q \rightarrow 1$, the assumption $\Gamma^2 = (\Gamma \otimes_{\mathcal{A}} \Gamma)/\mathcal{K}$ means that the higher order calculus $\Gamma^\wedge$ is some sense nearer to the corresponding construction in the classical case. Moreover, this assumption is essential in order to prove the following

THEOREM 5.2. *Suppose that g is an invariant metric. There is exactly one Levi-Civita connection on Γ with respect to g .*

PROOF : The proof is similar to that of Theorem 4.2. In Lemma 2.4 we proved that all invariant metrics have the form $g(\eta_{ij} \otimes \eta_{kl}) = (\alpha B_{14} B_{23} C_1^t C_3^t + \beta B_{12} B_{34} \hat{R}_{23} C_1^t C_3^t + \gamma B_{12} B_{34} C_1^t C_3^t)_{ijkl}$ with $\alpha + p\beta + \mathfrak{s}\gamma \neq 0$, $\alpha \neq q\beta$, $\alpha \neq -q^{-1}\beta$. By Lemma 5.1, the dimension of the vector space $\text{Mor}(u^c \otimes u, u^c \otimes u \otimes u^c \otimes u)$ is at most 15. A closer investigation of the proof of Lemma 5.1 shows that the following 15 morphisms $A_k = (A_{kij}^{mnrs})$ generate the vector space. (In case $N \geq 5$ they form a basis of this space.)

$$\begin{aligned} A_1 &= B_1^t B_3^t C_{12} C_{34} B_{12} C_1^t, & A_2 &= B_1^t B_3^t \hat{R}_{23}^{-1} C_{12} C_{34} B_{12} C_1^t, & A_3 &= B_1^t B_3^t C_{23} C_{14} B_{12} C_1^t, \\ A_4 &= B_3^t C_{34}, & A_5 &= B_1^t B_3^t \hat{R}_{12} C_{34} C_1^t, & A_6 &= B_1^t C_{12}, \\ A_7 &= B_1^t B_3^t C_{12} \hat{R}_{34} C_1^t, & A_8 &= B_3^t C_{23}, & A_9 &= B_1^t B_3^t C_{23} \hat{R}_{14} C_1^t, \\ A_{10} &= B_1^t B_3^t \hat{R}_{12} C_{23} C_1^t, & A_{11} &= B_1^t B_3^t \hat{R}_{12} C_{23} \hat{R}_{14}^{-1} C_1^t, & A_{12} &= B_3^t \hat{R}_{23}^{-1} C_{34}, \\ A_{13} &= B_1^t B_3^t \hat{R}_{23}^{-1} C_{34} \hat{R}_{12} C_1^t, & A_{14} &= B_1^t B_3^t \hat{R}_{12} \hat{R}_{23} C_{34} C_1^t, & A_{15} &= B_1^t B_3^t \hat{R}_{12} \hat{R}_{23} C_{34} \hat{R}_{12}^{-1} C_1^t. \end{aligned}$$

Therefore, by Lemma 3.2, we can make the following ansatz for our Levi-Civita connection ∇ :

$$\nabla(\eta_{ij}) = \sum_{k=1}^{15} \lambda_k A_{kij}^{mnrs} \eta_{mn} \otimes \eta_{rs}, \quad \lambda_k \in \mathbb{C}.$$

The condition for the vanishing torsion takes the form

$$(\sigma - \text{id})(\sigma - q^N \text{id})(\sigma - q^{-N} \text{id})(\eta \otimes \eta_{ij} + \eta_{ij} \otimes \eta - \sum_{k=1}^{15} \lambda_k A_{kij}^{mnrs} \eta_{mn} \otimes \eta_{rs}) = 0.$$

This leads to the equations

$$\begin{aligned} \lambda_4 &= \lambda_6 - Q\lambda_{15}, \quad \lambda_5 = \lambda_7 - Q\lambda_{14}, \quad \lambda_8 = pQ\lambda_6 + p\lambda_{15} - pQ, \quad \lambda_9 = pQ\lambda_7 + p\lambda_{14}, \\ \lambda_{11} - q^{-N} p\lambda_{10} &= Q(q^{-N} p\lambda_6 + q^{-N} (-q^{-N} p^2 + 1)\lambda_7 + \lambda_{15} - q^{-N} p - \frac{1}{p^2 + q^N} (p\lambda_6 + \lambda_7 + p\lambda_{14} + p^2\lambda_{15} - p)), \\ \lambda_{13} &= -Q\lambda_7 + q^N p^{-1} \lambda_{12} - q^N Q\lambda_{15} + \frac{q^N}{p^2 + q^N} (pQ\lambda_6 + Q\lambda_7 + pQ\lambda_{14} + p^2 Q\lambda_{15} - pQ). \end{aligned}$$

By Lemma 3.4, the connection is compatible with the metric if and only if the following equations are satisfied:

$$\begin{aligned} \lambda_6 \beta + \lambda_7 \alpha &= 0, \\ \lambda_6 (\alpha - Q\beta) + \lambda_7 \beta &= 0, \\ \lambda_1 (\alpha + p\beta + \mathfrak{s}\gamma) + \lambda_2 p^{-1} \gamma + \lambda_3 (Q\beta + \gamma) &= 0, \\ \lambda_2 \alpha + \lambda_3 \beta + \lambda_5 (\alpha + p\beta + \mathfrak{s}\gamma) + \lambda_9 (Q\beta + \gamma) + \lambda_{10} \gamma + \lambda_{13} p^{-1} \gamma + \lambda_{14} p (Q\beta + \gamma) &= 0, \\ \lambda_2 \beta + \lambda_3 (\alpha - Q\beta) + \lambda_4 (\alpha + p\beta + \mathfrak{s}\gamma) + \lambda_8 (Q\beta + \gamma) + \lambda_{11} \gamma + \lambda_{12} p^{-1} \gamma + \lambda_{15} p (Q\beta + \gamma) &= 0, \\ \lambda_8 (\alpha - Q\beta) + \lambda_{11} p\beta + \lambda_{12} \beta + \lambda_{15} p (\alpha - Q\beta) &= 0, \\ \lambda_8 \beta + \lambda_{10} p\beta + \lambda_{12} \alpha + \lambda_{14} p (\alpha - Q\beta) &= 0, \\ \lambda_9 (\alpha - Q\beta) + \lambda_{11} p\alpha + \lambda_{13} \beta + \lambda_{15} p\beta &= 0, \\ \lambda_9 \beta + \lambda_{10} p\alpha + \lambda_{13} \alpha + \lambda_{14} p\beta &= 0. \end{aligned}$$

Set $\tilde{p} = (q^N + 1)^{-1} \alpha_1^{-1} (p - p^{-3} q^{3N})$. Some computations show that the above system of equations admits a unique solution

$$\begin{aligned} 2\lambda_4 &= -Q^2(1 + \tilde{p}\beta), & 2\lambda_5 &= -Q^2\tilde{p}\alpha, & \lambda_6 &= \lambda_7 = 0, \\ 2\lambda_8 &= pQ(-1 + \tilde{p}\beta), & 2\lambda_9 &= pQ\tilde{p}\alpha, & 2\lambda_{10} &= Q(1 - \tilde{p}\beta), \\ 2\lambda_{11} &= -Q\tilde{p}(\alpha - Q\beta), & 2\lambda_{12} &= -pQ\tilde{p}(\alpha - Q\beta), & 2\lambda_{13} &= -pQ(1 + \tilde{p}\beta), \\ 2\lambda_{14} &= Q\tilde{p}\alpha, & 2\lambda_{15} &= Q(1 + \tilde{p}\beta), \end{aligned} \quad (23)$$

$$\lambda_3 = \frac{Q^2\alpha_0\alpha}{2\alpha_1\alpha_2} + Q\tilde{p}\gamma, \quad \lambda_2 = -\frac{Q^2\alpha_0\beta}{2\alpha_1\alpha_2} + \frac{Q^2}{2}\tilde{p}\alpha_0 - Qp\tilde{p}(Q\beta + \gamma),$$

$$\lambda_1 = \frac{Q^2}{2} \left(\frac{-Q\alpha\beta - \alpha\gamma + p^{-1}\beta\gamma}{\alpha_1\alpha_2} - p^{-1}\tilde{p}\gamma \right).$$

From the preceding considerations it is clear that the corresponding connection ∇ is indeed a Levi-Civita connection for g . $\blacksquare$

In order to examine the classical limit, we have to rewrite the Levi-Civita connection from Theorem 5.2 in terms of the standard basis. Using the projections P_0 , P_1 and P_2 defined in Section 2 some straightforward computations yield the formulas

$$\begin{aligned} \nabla(\eta) &= \frac{Q\alpha_0}{2\alpha_1} (pq^{-N} + p^{-1}q^N)(p^2q^{-N} - p^{-2}q^N) B_m^{ta} C^{md} P_1\eta_{ab} \otimes P_1\eta_{bd} + \\ &+ \frac{Q\alpha_0}{2} \left(\frac{Q\mathfrak{s}}{\alpha_2} + \tilde{p}(1 - q^N)(1 - q^N p^{-2}) \right) B_m^{ta} C^{md} P_2\eta_{ab} \otimes P_2\eta_{bd}, \end{aligned}$$

$$\begin{aligned} \nabla(P_1\eta_{ij}) &= \frac{Q(pq^{-N} + p^{-1}q^N)}{2} \left(\frac{p^{-2}q^N - p^2q^{-N}}{\mathfrak{s}} (\eta \otimes P_1\eta_{ij} + P_1\eta_{ij} \otimes \eta) - \right. \\ &- (p^2q^{-N} + 1)P_{1ij}^{kl} P_1\eta_{km} \otimes P_1\eta_{ml} + (p^{-2}q^{2N} - p^2q^{-N})P_{1ij}^{kl} P_2\eta_{km} \otimes P_1\eta_{ml} - \\ &- p\tilde{p}\alpha_1(1 + q^{-N})P_{1ij}^{kl} P_1\eta_{km} \otimes P_2\eta_{ml} + \tilde{p}\alpha_1(p^{-1}q^N - p)P_{1ij}^{kl} P_2\eta_{km} \otimes P_2\eta_{ml} \Big), \end{aligned}$$

$$\begin{aligned} \nabla(P_2\eta_{ij}) &= \frac{Q(pq^{-N} + p^{-1}q^N)}{2} \left(\frac{q^{-N} - q^N}{\mathfrak{s}} \eta \otimes P_2\eta_{ij} + \frac{(q^N - 1)(1 - p^{-2}q^N)\tilde{p}\alpha_2 - Q\mathfrak{s}}{(pq^{-N} + p^{-1}q^N)\mathfrak{s}} P_2\eta_{ij} \otimes \eta - \right. \\ &- (q^N + p^2q^{-2N})P_{2ij}^{kl} P_1\eta_{km} \otimes P_2\eta_{ml} + (1 - q^N)P_{2ij}^{kl} P_2\eta_{km} \otimes P_2\eta_{ml} + \\ &+ p\tilde{p}\alpha_2(1 + q^{-N})P_{2ij}^{kl} P_1\eta_{km} \otimes P_1\eta_{ml} + \tilde{p}\alpha_2(p - p^{-1}q^N)P_{2ij}^{kl} P_2\eta_{km} \otimes P_1\eta_{ml} \Big). \end{aligned}$$

From these formulas it follows that the classical limits of the Levi-Civita connections ∇ for both calculi Γ_+ and Γ_- exist. For the subspaces Υ^1 of 1-forms of Γ_+ which corresponds to the classical differential calculus (see [10]) we obtain

$$\begin{aligned} \nabla^{\text{cl}}(\bar{\omega}_{ij}^1) &= \lim_{q \rightarrow 1} \nabla(\omega_{ij}^1) = -\epsilon \frac{N - 2\epsilon}{N - \epsilon} (\bar{\omega}^0 \otimes \bar{\omega}_{ij}^1 + \bar{\omega}_{ij}^1 \otimes \bar{\omega}^0) - \vec{P}_{1ij}^{kl} \bar{\omega}_{km}^1 \otimes \bar{\omega}_{ml}^1 - \\ &- \frac{N - 4\epsilon}{N - 2\epsilon} \vec{P}_{1ij}^{kl} (\bar{\omega}_{km}^1 \otimes \bar{\omega}_{ml}^2 + \bar{\omega}_{km}^2 \otimes \bar{\omega}_{ml}^1 + \bar{\omega}_{km}^2 \otimes \bar{\omega}_{ml}^2). \end{aligned}$$

A The Rosso form of $\mathcal{U}_q(sl(N))$

In Section 2 we defined invariant metrics and we have seen that such metrics are not uniquely determined. On the other hand, Rosso showed in [17] that there is a unique ad-invariant bilinear form for the quantum universal enveloping algebra $\mathcal{U}_q(\mathfrak{g})$ for a simple Lie algebra $\mathfrak{g}$. In this appendix we define such a form adapted to the preceding considerations and we compute the corresponding ad-invariant metric on the quantum Lie algebra $\mathcal{X}$ of the FODC $\Gamma_{+,z}$.

Let q be a complex number, $q \neq 0$, $q^k \neq 1$ for all $k \in \mathbb{N}$ and let (a_{ij}) be the Cartan-matrix for $sl(N)$. Let $(\mathcal{U}_q(sl(N)), \Delta, \kappa, \varepsilon)$ be the Hopf algebra over $\mathbb{C}$ generated by the set of elements $\{E_i, F_i, K_i, K_i^{-1}, \tilde{K}_i, \tilde{K}_N, \tilde{K}_i^{-1}, \tilde{K}_N^{-1} \mid i = 1, \dots, N-1\}$ with relations

$$\begin{aligned} K_i K_j &= K_j K_i, & K_i K_i^{-1} &= K_i^{-1} K_i = 1, & K_i \tilde{K}_n &= \tilde{K}_n K_i, \\ \tilde{K}_m \tilde{K}_n &= \tilde{K}_n \tilde{K}_m, & \tilde{K}_n \tilde{K}_n^{-1} &= \tilde{K}_n^{-1} \tilde{K}_n = 1, & \tilde{K}_i^{-1} \tilde{K}_{i+1} &= K_i^2, \\ K_i E_j &= q^{a_{ij}} E_j K_i, & K_i F_j &= q^{-a_{ij}} F_j K_i \\ \tilde{K}_n E_j &= q^{2(\delta_{n,j+1} - \delta_{nj})} E_j \tilde{K}_n & \tilde{K}_n F_j &= q^{2(\delta_{nj} - \delta_{n,j+1})} F_j \tilde{K}_n, \\ E_i F_j - q^{-a_{ij}} F_j E_i &= \delta_{ij} \frac{K_i^2 - 1}{q^2 - 1}, \end{aligned} \tag{24}$$

$$\begin{aligned} \sum_{k=0}^{1-a_{ij}} (-1)^k b_{-a_{ij},k} E_i^k E_j E_i^{1-a_{ij}-k} &= 0 \quad (i \neq j), \\ \sum_{k=0}^{1-a_{ij}} (-1)^k b_{-a_{ij},k} F_i^k F_j F_i^{1-a_{ij}-k} &= 0 \quad (i \neq j) \end{aligned}$$

and coproduct Δ , antipode κ , counit ε defined by

$$\begin{aligned} \Delta(E_i) &= E_i \otimes K_i + 1 \otimes E_i, & \Delta(F_i) &= F_i \otimes K_i + 1 \otimes F_i, \\ \Delta(K_i) &= K_i \otimes K_i, & \Delta(\tilde{K}_n) &= \tilde{K}_n \otimes \tilde{K}_n, \\ \kappa(E_i) &= -q^2 K_i^{-1} E_i, & \kappa(F_i) &= -F_i K_i^{-1}, & \kappa(K_i) &= K_i^{-1}, & \kappa(\tilde{K}_n) &= \tilde{K}_n^{-1}, \\ \varepsilon(E_i) &= \varepsilon(F_i) = 0, & \varepsilon(K_i) &= \varepsilon(\tilde{K}_n) = 1 \end{aligned} \tag{25}$$

for all $i, j = 1, \dots, N-1$ and all $m, n = 1, \dots, N$. The constants in (24) are $b_{0,0} = b_{0,1} = b_{1,0} = b_{1,2} = 1$, $b_{1,1} = q + q^{-1}$. A realization of the Hopf algebra $\mathcal{U}_q(sl(N))$ in terms of the L -functionals is obtained by setting

$$E_i = Q^{-1} l^{-i} l^{+i}_{i+1}, \quad F_i = -q^{-1} Q^{-1} l^{-i+1} l^{+i+1}_{i+1}, \quad K_i = l^{-i} l^{+i+1}_{i+1}, \quad \tilde{K}_i = (l^{+i}_i)^2. \tag{26}$$

This Hopf algebra is $\mathbb{Z}^{N-1}$ -graduated with grading ∂ given by $\partial E_i := \alpha_i$, $\partial F_i = -\alpha_i$, $\partial K_i = 0$, $\partial \tilde{K}_i = 0$. Let $\langle \cdot, \cdot \rangle$ denote the symmetric bilinear form such that $\langle \alpha_i, \alpha_j \rangle = a_{ij}$.

It is well-known that all elements of $\mathcal{U}_q(sl(N))$ can be written as finite linear combinations of terms of the form FKE , where F and E are finite products of elements F_i and E_i , respectively, and K is a product of the generators K_i , K_i^{-1} , $\tilde{K}_i$ and $\tilde{K}_i^{-1}$.

Let us recall that a map $(\cdot, \cdot) : \mathcal{B} \times \mathcal{B} \rightarrow \mathbb{C}$ for a Hopf algebra $\mathcal{B}$ is called ad-invariant if

$$(\text{ad}_R \xi_{(1)}(\zeta_1), \text{ad}_R \xi_{(2)}(\zeta_2)) = \varepsilon(\xi)(\zeta_1, \zeta_2) \quad \text{for } \xi, \zeta_1, \zeta_2 \in \mathcal{B}. \quad (27)$$

Then one can prove the following proposition which is essentially Rosso's result adapted to the present setting. Note that in [17] the algebra $\mathcal{U}_q(\mathfrak{sl}(N))$ is different from ours and the left adjoint action is used.

PROPOSITION A.1. *There is a unique ad-invariant bilinear map $(\cdot, \cdot) : \mathcal{U}_q(\mathfrak{sl}(N)) \times \mathcal{U}_q(\mathfrak{sl}(N)) \rightarrow \mathbb{C}$ such that $(FKE, F'K'E') = (F, E')(K, K')(E, F')$, $(K, K') = (K', K)$, $(KK', K'') = (K, K'')(K', K'')$, $(K_i, K_j) = q^{-a_{ij}/2}$, $(\tilde{K}_i, K_j) = q^{\delta_{ij} - \delta_{i,j+1}}$ and $(\tilde{K}_i, \tilde{K}_j) = q^{-2\delta_{ij}}$.*

PROOF : The proof is similar to that of Theorem 6 in [17]. We omit the details. However, we want to stress that we deal with another adjoint action and with different commutation relations of the generators of $\mathcal{U}_q(\mathfrak{sl}(N))$. $\blacksquare$

In what follows we use the abbreviations $E_{i,i+1} := E_i$, $F_{i+1,i} := F_i$, $E_{i,j+1} := E_{i+1,j+1}E_i - q^{-1}E_iE_{i+1,j+1}$, $F_{j+1,i} := F_iF_{j+1,i+1} - q^{-1}F_{j+1,i+1}F_i$ for $i < j$. Using (27) and Proposition A.1 the bilinear form $(\cdot, \cdot)$ for these elements of $\mathcal{U}_q(\mathfrak{sl}(N))$ can be computed. The result is given by the formulas

$$\begin{aligned} (E_i, F_j) &= -q^{-1}Q^{-1}\delta_{ij}, & (E_{ij}, F_{kl}) &= -q^{2i-2j+1}Q^{-1}\delta_{il}\delta_{jk}, \\ (F_i, E_j) &= -qQ^{-1}\delta_{ij}, & (F_{ij}, E_{kl}) &= -qQ^{-1}\delta_{il}\delta_{jk}. \end{aligned}$$

From equations (2) and (26) the generators χ_{ij} of the quantum Lie algebra $\mathcal{X}$ of the FODC $\Gamma_{+,z}$ can be expressed in terms of the elements E_{ij} , F_{ji} , K_i and $\tilde{K}_i$:

$$\begin{aligned} \chi_{ij} &= q^{-1}QF_{ji}\tilde{K}_i + q^{-1}Q^2 \sum_{r < i} F_{jr}\tilde{K}_rE_{ri}, \quad (i < j) \\ \chi_{ij} &= q^{-1}Q\tilde{K}_jE_{ji} + q^{-1}Q^2 \sum_{r < j} F_{jr}\tilde{K}_rE_{ri}, \quad (i > j) \\ \chi_{ii} &= q^{-2}\tilde{K}_i - q^{-1}Q \sum_{m=1}^{i-1} q^{2m-2i}\tilde{K}_m + q^{-2}Q^2 \sum_{n < i} F_{in}\tilde{K}_nE_{ni} - \\ &\quad - q^{-1}Q^3 \sum_{m=1}^{i-1} \sum_{n=1}^{m-1} q^{2m-2i}F_{mn}\tilde{K}_nE_{nm} - q^{-2i}\varepsilon. \end{aligned}$$

Combining both sets of the preceding formulas it follows that the bilinear form in Proposition A.1 takes the following form on the generators χ_{ij} of the quantum Lie algebra:

$$(\chi_{ij}, \chi_{kl}) = -q^{-2i-1}Q\delta_{il}\delta_{jk} + q^{-2i-2j}Q^2\delta_{ij}\delta_{kl}. \quad (28)$$

This is precisely the dual metric g^* described by formula (10) for the parameter values $\alpha = -qQ^{-1}$, $\beta = -q^{2N+2}$ (cf. (10)).

B Another way to Levi-Civita connections

The results in Theorem 4.2 and 5.2 indicate that the concepts defined above are useful. Nevertheless, there are various other possibilities to define a metric and the compatibility of a metric with a connection. Here we pick out one of these possibilities which is similar to the classical case.

Let Γ denote one of the first order differential calculi $\Gamma_{\pm, z}$ of $SL_q(N)$ defined in Section 1. Suppose that q is real. Then Γ is a $*$ -calculus for the Hopf- $*$ -algebra $\mathcal{A} = SU_q(N)$. The involutions of $\mathcal{A}$ and of Γ are given by $(u_i^j)^* = \kappa(u_i^j)$ and $\eta_{ij}^* = -\eta_{ji}$, respectively.

We call a map $g : \Gamma \times \Gamma \rightarrow \mathcal{A}$ a **metric** if

$$g(a\xi, b\zeta) = ag(\xi, \zeta)b^* \quad \text{for all } a, b \in \mathcal{A}, \quad \xi, \zeta \in \Gamma.$$

We say a (left) connection ∇ on Γ is compatible with the metric g if

$$g(\nabla(\xi), \zeta) + g(\xi, \nabla(\zeta)) = dg(\xi, \zeta) \quad \text{for all } \xi, \zeta \in \Gamma.$$

This is well defined because of

$$\begin{aligned} g(\nabla(a\xi), b\zeta) + g(a\xi, \nabla(b\zeta)) - dg(a\xi, b\zeta) &= d(a)g(\xi, \zeta)b^* + ag(\nabla(\xi), \zeta)b^* + ag(\xi, \zeta)d(b^*) + \\ &+ ag(\xi, \nabla(\zeta))b^* - d(a)g(\xi, \zeta)b^* - adg(\xi, \zeta)b^* - ag(\xi, \zeta)d(b^*) = a(g(\nabla(\xi), \zeta) + g(\xi, \nabla(\zeta)) - dg(\xi, \zeta))b^*. \end{aligned}$$

A bicovariant left connection ∇ is called Levi-Civita connection in respect to the metric g , if ∇ has vanishing torsion and if it is compatible with the metric g . As above, such a Levi-Civita connection is uniquely determined by its values on the generators $\eta_{ij} \in \text{inv}\Gamma$.

Let g' be a metric from Lemma 2.3 with $\alpha, \beta \in \mathbb{R}$ and let g defined by $g(\xi, \zeta) = g'(\xi \otimes \zeta^*)$ with $\xi \in \Gamma, \zeta \in \text{inv}\Gamma$. Then we have $g(\eta_{ij}, \eta_{kl}) = q^{2j}\alpha\delta_{ik}\delta_{jl} + \beta\delta_{ij}\delta_{kl}$, $\alpha \neq 0$, $\alpha + \mathfrak{s}\beta \neq 0$. Let us assume that $\nabla(\eta_{ij}) = \sum_{k=1}^6 \lambda_k A_{kij}^{mnr} \eta_{mn} \otimes \eta_{rs}$ as in the proof of Theorem 4.2. In order to simplify the calculation, let λ_k be real.

The torsion of ∇ has to vanish and so we get (as in Theorem 4.2)

$$Q\lambda_3 = Q\lambda_4 = q\lambda_5 - (Q^2 + 1)\lambda_6 + Q.$$

The compatibility with the metric g gives only one equation

$$\lambda_2\alpha - \lambda_3(\alpha + \mathfrak{s}\beta) - \lambda_5\beta - q^{-1}\lambda_6\beta = 0.$$

Hence there is a three parameter family of Levi-Civita connections. Moreover, we have

$$\nabla(\eta) = (\lambda_2\mathfrak{s} + \lambda_5 + q^{-1}\lambda_6)q^{-2m}\eta_{mn} \otimes \eta_{nm} + (\lambda_1\mathfrak{s} + \lambda_3 + \lambda_4 + q^{2N}Q\lambda_6)\eta \otimes \eta$$

for the bi-invariant element $\eta \in \Gamma$. That is, even if we require in addition that $\nabla(\eta) = \lambda\eta \otimes \eta$ for some $\lambda \in \mathbb{C}$ we do not get a **unique** Levi-Civita connection.

References

- [1] Bresser, K., Dimakis, A., Müller-Hoissen, F. and Sitarz, A.: Noncommutative Geometry of Finite Groups. GOET-TP 95/95 (1995)
- [2] Brzezinski, T. and Majid, S.: Quantum Group Gauge Theory on Quantum Spaces. Commun. Math. Phys. **157**, 591–638 (1993)
- [3] Carow-Watamura, U., Schlieker, M., Watamura, S. and Weich, W.: Bicovariant Differential Calculus on Quantum Groups $SU_q(N)$ and $SO_q(N)$. Commun. Math. Phys. **142**, 605–641 (1991)

- [4] Connes, A.: *Non-Commutative Geometry*. Academic Press San Diego (1995)
- [5] Cuntz, J. and Quillen, D.: Algebra Extensions and Nonsingularity. *Journal of the American Mathematical Society* **8**(2), 251–289 (1995)
- [6] Drinfeld, V.G.: Quantum groups. In *Proceedings ICM 1986*, pages 798–820. Amer. Math. Soc. (1987)
- [7] Dubois-Violette, M. and Michor, W.: Connections on central bimodules. LPTHE-Orsay 94/100 (1995)
- [8] Faddeev, L.D., Reshetikhin, N.Yu. and Takhtajan, L.A.: Quantization of Lie Groups and Lie Algebras. *Algebra and Analysis* **1**, 178–206 (1987)
- [9] Georgelin, Y., Madore, J., Masson, T. and Mourad, J.: On the non-commutative Riemannian geometry of $GL_q(N)$. LPTHE-Orsay 95/51 (1995)
- [10] Heckenberger, I., Schmüdgen, K. and Schüler, A.: Classical limits of bicovariant differential calculi on the quantum groups $SL_q(N)$, $O_q(N)$ and $Sp_q(N)$. To appear.
- [11] Jimbo, M.: A q -analog of $\mathcal{U}(gl(N+1))$ Hecke algebras, and the Yang-Baxter equation. *Lett. Math. Phys.* **11**, 247–252 (1986)
- [12] Kassel, C.: *Quantum groups*. Grad. Texts in Math. Springer-Verlag Heidelberg (1995)
- [13] Lusztig, G.: Quantum deformations of certain simple modules over enveloping algebras. *Adv. Math.* **70**, 237–249 (1988)
- [14] Majid, S.: Quantum and Braided Lie Algebras. *J. Geom. Phys.* **13**, 307–356 (1994)
- [15] Majid, S.: *Foundations of Quantum Group Theory*. Cambridge University Press (1995)
- [16] Mourad, J.: Linear Connections in non-commutative geometry. *Class. Quantum Grav.* **12**, 965–974 (1995)
- [17] Rosso, M.: Analogues de la forme de Killing et du théorème d’Harish-Chandra pour les groupes quantiques. *Ann. scient. Éc. Norm. Sup.* **23**, 445–467 (1990)
- [18] Schmüdgen, K. and Schüler, A.: Classification of Bicovariant Differential Calculi on Quantum Groups of Type A, B, C and D. *Commun. Math. Phys.* **167**, 635–670 (1995)
- [19] Schmüdgen, K. and Schüler, A.: Classification of Bicovariant Differential Calculi on Quantum Groups. *Commun. Math. Phys.* **170**, 315–335 (1995)
- [20] Woronowicz, S.L.: Differential Calculus on Quantum Matrix Pseudogroups (Quantum Groups). *Commun. Math. Phys.* **122**, 125–170 (1989)